

GEODESIC RAY TRANSFORM WITH MATRIX WEIGHTS FOR PIECEWISE CONSTANT FUNCTIONS

JOONAS ILMAVIRTA AND JESSE RAILO

ABSTRACT. We show injectivity of the geodesic X-ray transform on piecewise constant functions when the transform is weighted by a continuous matrix weight. The manifold is assumed to be compact and nontrapping of any dimension, and in dimension three and higher we assume a foliation condition. We make no assumption regarding conjugate points or differentiability of the weight. This extends recent results for unweighted transforms.

1. INTRODUCTION

This article studies the weighted geodesic X-ray transform with injective matrix weights on nontrapping Riemannian manifolds with strictly convex boundary. This operator arises in many applications, and one of the basic questions is if the weighted line integrals over all maximal geodesics determine an unknown function. We show an injectivity result for a class of piecewise constant functions under the assumptions that the manifold admits a strictly convex function and the weight depends continuously on its coordinates on the unit sphere bundle. In two dimensions, the result follows for nontrapping manifolds with strictly convex boundary.

Let (M, g) be a compact nontrapping Riemannian manifold with strictly convex boundary. We denote by SM the unit sphere bundle and by Γ the set of maximal unit speed geodesics. The all geodesics have finite length by assumption. We say that M is *nontrapping* if every geodesic in Γ has finite length.

The *geodesic X-ray transform with matrix weights* is defined as follows. Fix some integers $m, k \geq 1$ and denote the set of linear injections (monomorphisms) $\mathbb{C}^k \rightarrow \mathbb{C}^m$ by $\text{Mon}(\mathbb{C}^k, \mathbb{C}^m)$, which is a subset of the space $\text{Lin}(\mathbb{C}^k, \mathbb{C}^m)$ of all linear maps. If $k = m$, we have $\text{Mon}(\mathbb{C}^k, \mathbb{C}^m) = GL(\mathbb{C}, k)$, and for $m < k$ we have $\text{Mon}(\mathbb{C}^k, \mathbb{C}^m) = \emptyset$. The geodesic X-ray transform with weight $W \in C(SM, \text{Lin}(\mathbb{C}^k, \mathbb{C}^m))$ is defined so that it maps a function $f: M \rightarrow \mathbb{C}^k$ to $I_W f: \Gamma \rightarrow \mathbb{C}^m$

Date: Friday 11th January, 2019.

2010 Mathematics Subject Classification. 44A12, 65R32, 53A99.

Key words and phrases. Geodesic ray transform, matrix weight, integral geometry, inverse problems.

University of Jyväskylä, Department of Mathematics and Statistics.

defined by

$$(1.1) \quad I_W f(\gamma) = \int_0^\tau W(\gamma(t), \dot{\gamma}(t)) f(\gamma(t)) dt$$

for any maximal geodesic $\gamma: [0, \tau] \rightarrow M$ whenever the integral is defined.

Injectivity of I_W for smooth functions was established by Paternain, Salo, Uhlmann, and Zhou [PSUZ16] if $\dim(M) \geq 3$, (M, g) admits a smooth strictly convex function, and $W \in C^\infty(SM; GL(k, \mathbb{C}))$. The result in [PSUZ16] is based on the methods developed in the work of Uhlmann and Vasy [UV16]. In this paper we consider a special case of the matrix weighted X-ray transform for the piecewise constant vector-valued functions. We gain more flexibility on geometrical assumptions and the proof is considerably simpler, but at the expense of only having the result for a restricted class of functions. Injectivity was shown recently in the case of piecewise constant functions without weights by Ilmavirta, Lehtonen, and Salo [ILS18], and reconstruction was studied in [Leb19]. Our main theorem is the following.

Theorem 1.1. *Let (M, g) be a compact nontrapping Riemannian manifold with strictly convex smooth boundary and $W \in C(SM; \text{Mon}(\mathbb{C}^k, \mathbb{C}^m))$. Suppose that either*

(a) $\dim(M) = 2$, or

(b) $\dim(M) \geq 3$ and (M, g) admits a smooth strictly convex function.

If $f: M \rightarrow \mathbb{C}^k$ is a piecewise constant function and $I_W f = 0$, then $f \equiv 0$.

There is also a local version of theorem 1.1; see theorem 2.6.

Remark 1.2. The result can be generalized by replacing \mathbb{C}^k and \mathbb{C}^m with two Banach spaces and letting W be an invertible linear map depending continuously on the coordinates on the sphere bundle SM .

Remark 1.3. The functions $f: M \rightarrow \mathbb{C}^k$ are vector-valued in the sense that they are sections of the trivial bundle $M \times \mathbb{C}^k$. We do not study geodesic X-ray tomography of vector fields or higher order tensor fields.

Theorem 1.1 generalizes results of [ILS18] for the matrix weighted X-ray transform similar to the one studied in [PSUZ16]. Our theorem holds if $\dim(M) \geq 2$, $W \in C(SM; GL(k, \mathbb{C}))$, and functions are piecewise constant in comparison to [PSUZ16] where it is assumed that $\dim(M) \geq 3$, $W \in C^\infty(SM; GL(k, \mathbb{C}))$ and functions are smooth. In the Euclidean space \mathbb{R}^n with $n \geq 3$ injectivity is known for $C^{1,\alpha}$ weights [Ilm16] but there is an example of non-injectivity for $W \in C^\alpha$ by Goncharov and Novikov [GN17]. Boman constructed an example of a smooth nonvanishing weight on the plane for which the weighted X-ray transform for smooth functions is non-injective [Bom93]. Theorem 1.1 shows that there are no such counterexamples for piecewise

Regularity	Dimension	$W = \text{Id}$	$W \in C^\infty$	$W \in C$
PWC	$= 2$	Yes. [ILS18]	Yes!	Yes!
L^2 or C^∞	$= 2$	Unknown.	No. [Bom93]	No. [Bom93]
PWC	≥ 3	Yes. [UV16]	Yes. [PSUZ16]	Yes!
L^2 or C^∞	≥ 3	Yes. [UV16]	Yes. [PSUZ16]	No. [GN17]

TABLE 1. Is the X-ray transform injective on manifolds that admit a strictly convex function? The answers in various different cases are summarized below. Here “PWC” stands for piecewise constant. A “Yes!” with an exclamation mark is a new result proven here. In two dimensions injectivity is known on simple manifolds, but the foliation condition does not imply simplicity.

constant functions. The known results — including the new ones obtained here — are summarized on Table 1.

We will prove Theorem 1.1 in Section 2. We remark that Theorem 1.1 is based on an generalization of [ILS18, Lemma 4.2] whereas the rest of the proof is almost identical to the one in [ILS18]. The method in [ILS18] relies on existence of a strictly convex foliation as in the works of Stefanov, Uhlmann, and Vasy [UV16, SUV17], but the method of proof is far simpler. For a further discussion on the foliation condition see [PSUZ16] and references therein.

The matrix weighted X-ray transform is related to recovering matrix valued connection from its parallel transport [FU01, Nov02a, PSU12, GPSU16, PSUZ16]. It has also applications in polarization tomography [Sha94, NS07, Hol13] and quantum state tomography [Ilm16].

One source of weights is pseudolinearization, a procedure where a nonlinear problem is reduced to a linear problem with weights depending on the unknown. For a more detailed description of the idea, first appearing in [SU98, SUV16] (see e.g. [IM19, Section 8]). Pseudolinearization also leads to an iterative inversion algorithm [Ilm16, SU98, SUV16].

A boundary reconstruction of the normal derivatives of a function from the broken ray transform reduces to a certain weighted geodesic ray transform on the boundary [Ilm14]. Some weights can be realized as attenuation, but we make no such assumptions on W . The attenuated X-ray transform (see e.g. [ABK97, Nov02b, SU11, AMU18]) is a well-known special case of the matrix weighted X-ray transform and it is the mathematical basis for the medical imaging method SPECT (see e.g. the survey [Fin03]).

Acknowledgements. J.I. was supported by the Academy of Finland (decision 295853). J.R. was supported by the Academy of Finland (Centre of Excellence in Inverse Problems Research at the University




FIGURE 1. An example of a regular tiling and the tangent space at the point x . The simplices touching x have their corresponding tangent cones on $T_x M$. See Definition 2.1.

of Jyväskylä in 2017, Centre of Excellence in Inverse Modelling and Imaging at the University of Helsinki in 2018). The authors are grateful to Jere Lehtonen and Mikko Salo for helpful discussions related to this work.

2. PROOF

2.1. Definitions. We follow the notation of [ILS18], and any details omitted here can be found there. We review the main concepts here in a somewhat informal manner. A regular m -simplex on a manifold M is a C^1 -smoothly embedded standard m -dimensional simplex. We denote $n = \dim(M)$. A regular tiling of a manifold is a collection of regular n -simplices which cover the manifold, whose interiors are disjoint, and whose boundaries align nicely. An example is given in Figure 1. A piecewise constant function is such that the values are constant in the interior of every simplex and zero on their boundaries. The geometry of corners of simplices is important for our argument, and we review the crucial definitions in more depth.

Definition 2.1 (Tangent cone). Let Δ be a regular m -simplex in M with $0 \leq m \leq n = \dim(M)$, and let $x \in \Delta$. Let $\Gamma = \Gamma(x, \Delta)$ be the set of all C^1 -curves starting at x and staying in Δ . The *tangent cone* of Δ at x is the set

$$(2.1) \quad C_x \Delta := \{ \dot{\gamma}(0) \mid \gamma \in \Gamma \}.$$

Definition 2.2 (Tangent function). Let $f: M \rightarrow \mathbb{C}^k$ be a piecewise constant function and $x \in M$ with respect to a regular tiling. Let $\Delta_1, \dots, \Delta_N$ be the simplices of the regular tiling that contain x . Denote

by $v_1, \dots, v_N \in \mathbb{C}^k$ the constant values of f in the interior of these simplices. The *tangent function* $T_x f: T_x M \rightarrow \mathbb{C}^k$ of f at x is defined so that for each $i \in \{1, \dots, N\}$ the function $T_x f$ takes the constant value v_i in the interior of the tangent cone $C_x \Delta_i$. The tangent function takes the value zero in $T_x M \setminus \bigcup_{i=1}^N \text{Int}(C_x \Delta_i)$.

We stress that the tangent function is not a derivative, as a piecewise constant function is typically not differentiable at the points of interest. Instead of linearizing the function, we linearize the geometry of the simplices and keep the constant values of the function.

2.2. Lemmas. In this subsection we recall a key lemma proved in [ILS18, Section 4] and use it to prove a new lemma.

Let M be a C^2 -smooth Riemannian surface with C^2 -boundary. Suppose the boundary ∂M is strictly convex at $x \in \partial M$. Let $\gamma_i, i = 1, 2$, be two unit speed C^1 -curves in M starting nontangentially at x so that $|\langle \dot{\gamma}_1(0), \dot{\gamma}_2(0) \rangle| < 1$.

Let $r > 0$ be such that $B(x, r) \subset M$ is split by the curves into three parts. Let A be the middle one. Let $\sigma_i, i = 1, 2$, be the curves on $T_x M$ with constant speed $\dot{\gamma}_i(0)$ respectively. Let S be the sector in $T_x M$ laying between σ_1 and σ_2 .

If $h > 0$ and if $v \in T_x M$ is an inward pointing unit vector, let the geodesic γ_v^h be constructed as follows: Take a unit vector w normal to v at x — which is unique up to sign — and let $w(h)$ be its parallel transport along the geodesic through v by distance h . Let γ_v^h be the maximal geodesic in the direction of $w(h)$. Let σ_v^h be the corresponding line $h\nu + w\mathbb{R}$ in $T_x M$. We denote by $\nu \in T_x \partial M$ the inward pointing unit vector of ∂M at x .

We restate a lemma from [ILS18] for convenience.

Lemma 2.3 ([ILS18, Lemma 4.1]). *Let M be a C^2 -smooth Riemannian surface with C^2 -boundary, which is strictly convex at $x \in \partial M$. There exists an open neighborhood U of ν such that for every $v \in U$ we have*

$$(2.2) \quad \lim_{h \rightarrow 0} \frac{1}{h} \int_{\gamma_v^h \cap A} ds = \int_{\sigma_v^h \cap S} ds.$$

Lemma 2.3 allows us to reduce the local problem of Theorem 1.1 into the corresponding local problem in the tangent space. We next state a generalization of [ILS18, Lemma 4.2] for the class of piecewise constant vector-valued functions with matrix weighted integrals.

Lemma 2.4. *Let M be a C^2 -smooth Riemannian surface with C^2 boundary, which is strictly convex at $x \in \partial M$. Let \tilde{M} be such an extension of M that $x \in \text{Int}(\tilde{M})$. Let $\Delta \subset \tilde{M}$ be a regular 2-simplex so that $C_x \Delta \cap T_x \partial M = \{0\}$. Let $W \in C(\tilde{M}; \text{Lin}(\mathbb{C}^k, \mathbb{C}^m))$ and $f: \tilde{M} \rightarrow \mathbb{C}^k$ be a piecewise constant function supported in Δ . Then there exists an*

open neighborhood U of ν such that for every $v \in U$ we have

$$(2.3) \quad \lim_{h \rightarrow 0} \frac{1}{h} \int_{\gamma_v^h} W(\gamma_v^h(s), \dot{\gamma}_v^h(s)) f(\gamma_v^h(s)) ds = W(x, v^\perp) \int_{\sigma_v^1} T_x f(\sigma_v^1(s)) ds.$$

Proof. By linearity we can assume that f is constant in Δ . A piecewise constant function is a linear combination of characteristic functions of interiors of simplices.

Fix $v \in U$ given by Lemma 2.3. Let $s_h \in \gamma_v^h$ be any maximizer of

$$(2.4) \quad s \mapsto \|W(\gamma_v^h(s), \dot{\gamma}_v^h(s)) - W(x, v^\perp)\|,$$

where — as throughout this proof — we use the operator norm of matrices. We have

$$(2.5) \quad \sup_{s \in \gamma_v^h} d((\gamma_v^h(s), \dot{\gamma}_v^h(s)), (x, v^\perp)) \rightarrow 0$$

as $h \rightarrow 0$, and so $(\gamma_v^h(s_h), \dot{\gamma}_v^h(s_h)) \rightarrow (x, v^\perp)$ as $h \rightarrow 0$. We have $\lim_{h \rightarrow 0} \frac{l(\gamma_v^h \cap \Delta)}{h} = |\sigma_v^1|$, and in particular the fraction $\frac{l(\gamma_v^h \cap \Delta)}{h}$ is uniformly bounded for all small $h > 0$.

We ready to compare the weighted integral on the left-hand side of (2.3) to the corresponding integral with the weight frozen to its limit value $W(x, v^\perp)$ (as $h \rightarrow 0$). Straightforward estimates give

$$(2.6) \quad \left\| \frac{1}{h} \int_{\gamma_v^h} (W(\gamma_v^h(s), \dot{\gamma}_v^h(s)) - W(x, v^\perp)) f ds \right\| \leq \frac{l(\gamma_v^h \cap \Delta)}{|h|} \sup_{s \in \gamma_v^h} \|f(s)\| \sup_{s \in \gamma_v^h} \|W(\gamma_v^h(s), \dot{\gamma}_v^h(s)) - W(x, v^\perp)\|.$$

As f is bounded, the quotient $\frac{l(\gamma_v^h \cap \Delta)}{h}$ is bounded, and the matrix norm tends to zero as $h \rightarrow 0$, the left-hand side of (2.6) tends to zero as well.

We may thus conclude that the limit on the left-hand side of (2.3) is the same as

$$(2.7) \quad W(x, v^\perp) \lim_{h \rightarrow 0} \frac{1}{h} \int_{\gamma_v^h} f(\gamma_v^h(s)) ds.$$

That is, the weight can be frozen to its limiting value. The function f is constant, so up to that constant the integral is just the length of the geodesic segment in Δ . Lemma 2.3 shows that the characteristic function of a simplex satisfies (2.3) in the absence of weight. This concludes the proof. \square

Suppose Σ is a hypersurface containing the point $x \in \text{Int}(M)$ and Σ is strictly convex in a neighbourhood of x . Let V be a small neighbourhood of x such that $V \setminus \Sigma$ consists of two open sets which are denoted by V_+ and V_- . We choose V_+ to be the one for which the boundary section $\partial V_+ \cap \Sigma$ is strictly convex. Next we state Lemma 2.5

that allows one to build a layer stripping argument that is used to prove Theorem 1.1.

Lemma 2.5. *Let M be a C^2 -smooth Riemannian manifold, $W \in C(M; \text{Mon}(\mathbb{C}^k, \mathbb{C}^m))$ and $f: M \rightarrow \mathbb{C}^k$ be a piecewise constant function. Fix $x \in \text{Int}(M)$ and let Σ be an $(n-1)$ -dimensional hypersurface through x . Suppose that V is a neighbourhood of x so that*

- V intersects only simplices containing x ,
- Σ is strictly convex in V ,
- $f|_{V_-} = 0$, and
- $I_W f = 0$ over every maximal geodesic in V having endpoints on Σ .

Then $f|_V = 0$.

Proof. The lemma follows from Lemma 2.4 using ideas developed in [ILS18, Lemma 5.1 and Lemma 6.2]. We summarize the idea briefly; details can be found in the cited paper.

Consider two dimensions first. Take a unit vector $v \in T_x M$ pointing towards V_+ and $h > 0$. Define the geodesic γ_v^h as above. The weighted integrals of f over these geodesics vanish by assumption. By Lemma 2.4 and injectivity of W everywhere on the sphere bundle, we find that $T_x f$ integrates to zero over σ_v^1 .

This argument reduces the X-ray tomography problem on M to the corresponding problem on $T_x M$. We have the freedom to vary the direction v , and any open set is sufficient. The Euclidean problem is unweighted and can be solved by explicit calculation; see [ILS18, Lemma 3.1]. The calculation is based on describing the direction v by a parameter and computing derivatives of high orders with respect to that parameter. That these derivatives determine the values of $T_x f$ in the cones uniquely boils down to the invertibility of a Vandermonde matrix.

In higher dimensions one can take a suitable two-dimensional surface through the point x and consider the problem on this submanifold. The above argument shows that f has to vanish on all simplices meeting x and this surface. There is sufficient freedom in choosing the surface so that all simplices are covered and thus the proof is complete. \square

2.3. Proof of Theorem 1.1. We are now ready to prove our main theorem. We begin with a local version.

Following [PSUZ16], we say that a subset $U \subset M$ has a strictly convex foliation if there is a strictly convex function $\phi: U \rightarrow \mathbb{R}$ so that the sets $\{x \in U; \phi(x) \geq c\}$ for all $c > \inf_U \phi$ are compact.

Theorem 2.6. *Let M be a C^2 -smooth Riemannian manifold with strictly convex boundary and $\dim(M) \geq 2$. Suppose that a subset $U \subset M$ has a strictly convex foliation. Let $W \in C(M; GL(k, \mathbb{C}))$ and $f: M \rightarrow \mathbb{C}^k$*

be a piecewise constant function. If $I_W f = 0$ for all geodesics in U , then $f|_U = 0$.

Proof. The proof is very similar to that of [ILS18, Theorem 5.3 and Theorem 6.4], and we only give an outline.

The set U can be foliated by strictly convex hypersurfaces, and we are interested at the times when the foliation meets a new simplex. It suffices to prove local injectivity in the neighborhood of a strictly convex boundary point — a point on the leaf of a foliation — whenever a new simplex is met. If the point meets only one new simplex, then one can use a sequence of short geodesics that pass the simplex and argue as in Lemma 2.4 to see that f has to vanish in the simplex. If the point meets more simplices, we are in the setting of Lemma 2.5, and we can conclude that the function vanishes on each new simplex. \square

Theorem 2.6 has also corollaries analogous to [ILS18, Corollaries 6.5–6.7] but we omit them here.

Proof of Theorem 1.1. Under the assumptions there is a global foliation: we may choose $U = M$ in Theorem 2.6. See [PSUZ16, Section 2] for details. The proof is complete. \square

REFERENCES

- [ABK97] È. V. Arbuzov, A. L. Bukhgeim, and S. G. Kazantsev. Two-dimensional problems in tomography, and the theory of A -analytic functions. In *Algebra, geometry, analysis and mathematical physics (Russian) (Novosibirsk, 1996)*, pages 6–20, 189. Izdat. Ross. Akad. Nauk Sib. Otd. Inst. Mat., Novosibirsk, 1997.
- [AMU18] Yernat M. Assylbekov, François Monard, and Gunther Uhlmann. Inversion formulas and range characterizations for the attenuated geodesic ray transform. *J. Math. Pures Appl. (9)*, 111:161–190, 2018.
- [Bom93] Jan Boman. An example of nonuniqueness for a generalized Radon transform. *J. Anal. Math.*, 61:395–401, 1993.
- [Fin03] David V. Finch. The attenuated x-ray transform: recent developments. 47:47–66, 2003.
- [FU01] David Finch and Gunther Uhlmann. The x-ray transform for a non-abelian connection in two dimensions. *Inverse Problems*, 17(4):695–701, 2001. Special issue to celebrate Pierre Sabatier’s 65th birthday (Montpellier, 2000).
- [GN17] Fedor Goncharov and Roman Novikov. A breakdown of injectivity for weighted ray transforms in multidimensions. 2017. Preprint, arXiv:1711.06163.
- [GPSU16] Colin Guillarmou, Gabriel P. Paternain, Mikko Salo, and Gunther Uhlmann. The X-ray transform for connections in negative curvature. *Comm. Math. Phys.*, 343(1):83–127, 2016.
- [Hol13] Sean Holman. Generic local uniqueness and stability in polarization tomography. *J. Geom. Anal.*, 23(1):229–269, 2013.
- [Ilm14] Joonas Ilmavirta. Boundary reconstruction for the broken ray transform. *Ann. Acad. Sci. Fenn. Math.*, 39(2):485–502, 2014.
- [Ilm16] Joonas Ilmavirta. Coherent quantum tomography. *SIAM J. Math. Anal.*, 48(5):3039–3064, 2016.

- [ILS18] Joonas Ilmavirta, Jere Lehtonen, and Mikko Salo. Geodesic x-ray tomography for piecewise constant functions on nontrapping manifolds. *Mathematical Proceedings of the Cambridge Philosophical Society*, page 1–13, 2018.
- [IM19] Joonas Ilmavirta and François Monard. Integral geometry on manifolds with boundary and applications. In Ronny Ramlau and Otmar Scherzer, editors, *The Radon Transform: The First 100 Years and Beyond*. de Gruyter, 2019. To appear; preprint at arXiv:1806.06088.
- [Leb19] Vadim Lebovici. Reconstruction of piecewise constant functions from X-ray data. 2019. Preprint, arXiv:1901.01909.
- [Nov02a] R. G. Novikov. On determination of a gauge field on \mathbb{R}^d from its non-abelian Radon transform along oriented straight lines. *J. Inst. Math. Jussieu*, 1(4):559–629, 2002.
- [Nov02b] Roman G. Novikov. An inversion formula for the attenuated X-ray transformation. *Ark. Mat.*, 40(1):145–167, 2002.
- [NS07] Roman Novikov and Vladimir Sharafutdinov. On the problem of polarization tomography. I. *Inverse Problems*, 23(3):1229–1257, 2007.
- [PSU12] Gabriel P. Paternain, Mikko Salo, and Gunther Uhlmann. The attenuated ray transform for connections and Higgs fields. *Geom. Funct. Anal.*, 22(5):1460–1489, 2012.
- [PSUZ16] Gabriel P. Paternain, Mikko Salo, Gunther Uhlmann, and Hanming Zhou. The geodesic x-ray transform with matrix weights. 2016. Preprint, arXiv:1605.07894v2.
- [Sha94] Vladimir A. Sharafutdinov. *Integral geometry of tensor fields*. Inverse and Ill-posed Problems Series. VSP, Utrecht, 1994.
- [SU98] Plamen Stefanov and Gunther Uhlmann. Rigidity for metrics with the same lengths of geodesics. *Math. Res. Lett.*, 5(1-2):83–96, 1998.
- [SU11] Mikko Salo and Gunther Uhlmann. The attenuated ray transform on simple surfaces. *J. Differential Geom.*, 88(1):161–187, 2011.
- [SUV16] Plamen Stefanov, Gunther Uhlmann, and Andras Vasy. Boundary rigidity with partial data. *J. Amer. Math. Soc.*, 29(2):299–332, 2016.
- [SUV17] Plamen Stefanov, Gunther Uhlmann, and András Vasy. Local and global boundary rigidity and the geodesic x-ray transform in the normal gauge. 2017. Preprint, arXiv:1702.03638.
- [UV16] Gunther Uhlmann and András Vasy. The inverse problem for the local geodesic ray transform. *Invent. Math.*, 205(1):83–120, 2016.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF JYVÄSKYLÄ,
P.O. BOX 35 (MAD) FI-40014 UNIVERSITY OF JYVÄSKYLÄ, FINLAND

E-mail address: `joonas.ilmavirta@jyu.fi`

E-mail address: `jesse.t.railo@jyu.fi`