

# SOLUTIONS OF NONLINEAR PDES IN THE SENSE OF AVERAGES

BERND KAWOHL, JUAN MANFREDI, AND MIKKO PARVIAINEN

ABSTRACT. We characterize  $p$ -harmonic functions including  $p = 1$  and  $p = \infty$  by using mean value properties extending classical results of Privaloff from the linear case  $p = 2$  to all  $p$ 's. We describe a class of random tug-of-war games whose value functions approach  $p$ -harmonic functions as the step goes to zero for the full range  $1 < p < \infty$ .

## 1. INTRODUCTION

In this article we study solutions to a class of non-linear equations that can be characterized by mean value properties. The quintessential example is the characterization of harmonic functions by the property

$$(1) \quad u(x) = \int_{B_\varepsilon(x)} u(y) dy.$$

Privaloff [24] proved that an upper semicontinuous function  $u$  is subharmonic if and only if

$$(2) \quad \limsup_{\varepsilon \rightarrow 0} \frac{1}{\varepsilon^2} \left[ \int_{B_\varepsilon(x)} u(y) dy - u(x) \right] \geq 0.$$

A similar statement for spherical means was obtained by Blasche [1].

If we replace the Laplace equation  $\Delta u = 0$  by a linear elliptic equation with constant coefficients  $Lu = \sum_{i,j} a_{ij} u_{x_i x_j} = 0$  then mean value formulas now hold for appropriate ellipsoids instead of balls. This is true also in the subelliptic case. See Chapter 5 in the recent book [3] and the paper [2] for updated discussion of mean value properties for solutions of linear equations.

We are interested in understanding mean value properties in the nonlinear case. We start by observing that in order to characterize continuous harmonic functions it is enough to ask that the mean value property (1) holds in an

---

2010 *Mathematics Subject Classification.* 35J92, 35C15, 35C20, 31B35, 35D40, 49L25, 91A15.

*Key words and phrases.* 1-harmonic, asymptotic expansion, dynamic programming principle, mean value theorem,  $p$ -harmonic, stochastic games, two-player zero-sum games, viscosity solutions.

asymptotic sense

$$(3) \quad u(x) = \int_{B_\varepsilon(x)} u(y) dy + o(\varepsilon^2) \text{ as } \varepsilon \rightarrow 0.$$

In fact, even a weaker *viscosity* notion suffices. An upper semicontinuous function  $u: \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}$  is *subharmonic* in  $\Omega$  if for every  $x \in \Omega$  and test function  $\phi \in C^2(\Omega)$  that touches  $u$  from above at  $x$  we have that

$$(4) \quad \phi(x) \leq \int_{B_\varepsilon(x)} \phi(y) dy + o(\varepsilon^2) \text{ as } \varepsilon \rightarrow 0.$$

Notice that the characterization (4) implies a simple proof of one half of Privaloff's characterization. Similarly, solutions to the  $p$ -Laplace equation are characterized by

$$(5) \quad u(x) = \frac{p-2}{2(p+n)} \left\{ \max_{\overline{B_\varepsilon(x)}} u + \min_{\overline{B_\varepsilon(x)}} u \right\} + \frac{n+2}{p+n} \int_{B_\varepsilon(x)} u(y) dy + o(\varepsilon^2)$$

in the viscosity sense, for  $p$  in the range  $1 < p \leq \infty$ . These facts are proven in [19]. That is, we have the analogue of Privaloff's characterization for  $p$ -subharmonic functions by replacing the regular solid average with the nonlinear average in (5) and using expansions in the viscosity sense. For a related evolution problem see [7] as well as [20], and for general discussion of  $p$ -Laplacian problems including  $p = 1$  and  $p = \infty$  see [13].

In [22], Peres and Sheffield showed that  $p$ -harmonic functions are limits of value functions of certain tug-of-war games with noise as the step size tends to zero. These games were modified in [21] so that their value function  $u_\varepsilon$ , with step size  $\varepsilon > 0$ , was uniquely defined, and satisfied a dynamic programming principle of the form

$$u_\varepsilon(x) = \frac{p-2}{2(p+n)} \left\{ \max_{\overline{B_\varepsilon(x)}} u_\varepsilon + \min_{\overline{B_\varepsilon(x)}} u_\varepsilon \right\} + \frac{n+2}{p+n} \int_{B_\varepsilon(x)} u_\varepsilon(y) dy,$$

when  $p \geq 2$ . See also [15, 16] and [23].

The objectives of this paper are to consider the limit case  $p = 1$ , to characterize the 1-harmonic functions in the spirit of (5), and to obtain a dynamic programming principle valid for all  $p > 1$  for the corresponding tug-of-war game. Theorem 9 below states that 1-harmonic subsolutions are characterized by

$$u(x) = \int_{B_\varepsilon^{\pi\nu_{\min}}} u(x+h) d\mathcal{L}^{n-1}(h) + o(\varepsilon^2),$$

where  $u(x + \varepsilon\nu_{\min}) = \min_{y \in \overline{B_\varepsilon(x)}} u(y)$ , and  $B_\varepsilon^{\pi\nu_{\min}}$  is the  $(n-1)$ -dimensional ball centered at zero in the hyperplane  $\pi_{\nu_{\min}}$ , which is perpendicular to  $\nu_{\min}$ . Both definitions are to be understood in the viscosity sense. In Theorems 11 and 13,

we extend this formula to the whole  $p$ -range by interpolating between the 1-Laplacian and the infinity Laplacian. Finally, we state the dynamic programming principle in Lemma 14.

In Section §2 we review the various definitions of viscosity solutions for the  $p$ -Laplacian. In Section §3 we study the limit case  $p = 1$ . The results of this section are used in §4, where we consider  $p$ -harmonic functions in the sense of averages for all  $p > 1$ . The corresponding asymptotic mean value characterization is derived in Section §5. Finally in §6 we describe a tug of war game whose value function satisfies an appropriate dynamic programming principle.

## 2. VISCOSITY SOLUTIONS OF THE NORMALIZED $p$ -LAPLACIAN

Let  $u$  be a real valued function of class  $C^2$  with non-vanishing gradient. For  $p \in [1, \infty)$  the normalized version of the  $p$ -Laplace operator acting on  $u$  is

$$\Delta_p^N u = \frac{1}{p} |\nabla u|^{2-p} \Delta_p u = \frac{1}{p} |\nabla u|^{2-p} \operatorname{div}(|\nabla u|^{p-2} \nabla u)$$

while for  $p = \infty$  we set

$$\Delta_\infty^N u = |\nabla u|^{-2} \Delta_\infty u = |\nabla u|^{-2} \langle D^2 u \nabla u, \nabla u \rangle$$

After a calculation we see that

$$\begin{aligned} \operatorname{div}(|\nabla u|^{p-2} \nabla u) &= |\nabla u|^{p-2} ((p-2) |\nabla u|^{-2} \langle D^2 u \nabla u, \nabla u \rangle + \Delta u) \\ &= |\nabla u|^{p-2} ((p-2) \Delta_\infty^N u + \Delta u) \\ (6) \quad &= |\nabla u|^{p-2} ((p-1) \Delta_\infty^N u + (\Delta u - \Delta_\infty^N u)) \\ &= |\nabla u|^{p-2} ((p-1) \Delta_\infty^N u + \Delta_1^N u), \end{aligned}$$

and thus

$$(7) \quad \Delta_p^N u = \frac{1}{p} \Delta_1^N u + \frac{1}{q} \Delta_\infty^N u,$$

where  $q$  is the Hölder conjugate of  $p$ ,  $1/p + 1/q = 1$ . Note that  $\Delta_2^N = (1/2) \Delta u$  and that

$$\Delta u = \Delta_1^N + \Delta_\infty^N.$$

As the name viscosity solution suggests, one of their origins lies in adding an artificial viscosity term  $\varepsilon \Delta u$  to a degenerate elliptic equation and sending  $\varepsilon$  to zero. For the normalized 1-Laplacian this amounts to studying equations like  $\varepsilon \Delta u_\varepsilon + \Delta_1^N u_\varepsilon = 0$  and this equation can be rewritten as  $(2\varepsilon + 1) \Delta_{p_\varepsilon}^N u_\varepsilon = 0$  with  $p_\varepsilon = 1 + \frac{\varepsilon}{1+\varepsilon}$ . Notice that  $p_\varepsilon \rightarrow 1$  as  $\varepsilon \rightarrow 0$ . In fact, it is known that a sequence of  $p$ -harmonic functions converges to a 1-harmonic function as  $p \rightarrow 1$ ; see [6] and also the discussion before Theorem 6. Moreover, a special 1-harmonic limit is chosen. It is a function of least gradient as pointed out by Juutinen in [10, Remark 3.3].

To deal with the case of non-smooth functions, we define  $\mathbb{F}_p(\eta, X)$  for  $\eta \in \mathbb{R}^n \setminus \{0\}$  and for symmetric  $n \times n$  matrices  $X$  as

$$\mathbb{F}_p(\eta, X) = \sum_{i,j}^n \left( \frac{1}{p} \delta_{ij} + \frac{(p-2)}{p} \frac{\eta_i \eta_j}{|\eta|^2} \right) X_{ij}$$

and for  $p = \infty$  as

$$\mathbb{F}_\infty(\eta, X) = \sum_{i,j}^n \left( \frac{\eta_i \eta_j}{|\eta|^2} \right) X_{ij},$$

so that we always have

$$\mathbb{F}_p(\eta, X) = \frac{1}{p} \mathbb{F}_1(\eta, X) + \frac{1}{q} \mathbb{F}_\infty(\eta, X).$$

These functions are used to define viscosity solutions to the nonhomogeneous problem for the  $p$ -Laplace operator in non-divergence form

$$(8) \quad \begin{aligned} \Delta_p^N u(x) &= \left[ \frac{1}{p} |\nabla u|^{2-p} \operatorname{div}(|\nabla u|^{p-2} \nabla u) \right] (x) \\ &= \mathbb{F}_p(\nabla u(x), D^2 u(x)) = g(x). \end{aligned}$$

To define a viscosity solution to the equation

$$(9) \quad \mathbb{F}_p(\nabla u, D^2 u) = g(x)$$

with  $g \in C(\bar{\Omega})$ ,  $g > 0$  (or  $g < 0$ ), we need to compute  $\mathbb{F}_p(\nabla \phi(x), D^2 \phi(x))$  for  $C^2$ -smooth test functions touching  $u$  from above or below at the test point  $x \in \Omega$ . Unfortunately, except for  $p = 2$ , the functions  $\mathbb{F}_p(\eta, X)$  are discontinuous when  $\eta = 0$ . There are several ways in the literature, see for example [6], [9], and [23], to resolve this difficulty.

- (1) We can modify our requirements when  $\nabla \phi(x) = 0$ ,
- (2) we can restrict the class of test functions so that  $\mathbb{F}_p(\nabla \phi(x), D^2 \phi(x))$  is uniquely defined also when  $\nabla \phi(x) = 0$  by  $\lim_{\eta \rightarrow 0} \mathbb{F}_p(\eta, D^2 \phi(x))$ , and
- (3) we can extend the domain of  $\mathbb{F}_p(\eta, X)$  by using semicontinuous extensions.

Let us start with the first approach.

**Definition 1.** *A continuous function  $v$  is a viscosity solution to the equation*

$$\mathbb{F}_p(\nabla v, D^2 v) = g(x)$$

*at  $x$ , if and only if every  $C^2$ -function  $\phi$  that touches  $v$  from below in  $x$  satisfies*

$$\mathbb{F}_p(\eta, D^2 \phi(x)) \leq g(x) \quad \begin{cases} \text{for } \eta = \nabla \phi(x) & \text{if } \nabla \phi(x) \neq 0, \\ \text{for some } \eta \in B_1(0) \setminus \{0\} & \text{if } \nabla \phi(x) = 0 \end{cases}$$

and every  $C^2$ -function  $\phi$  that touches  $u$  from above at  $x$  satisfies

$$\mathbb{F}_p(\eta, D^2\phi(x)) \geq g(x) \quad \begin{cases} \text{for } \eta = \nabla\phi(x) & \text{if } \nabla\phi(x) \neq 0, \\ \text{for some } \eta \in B_1(0) \setminus \{0\} & \text{if } \nabla\phi(x) = 0. \end{cases}$$

By saying that  $\phi$  touches  $u$  from below at  $x_0$ , we mean

- i)  $u(x_0) = \phi(x_0)$ ,
- ii)  $u(x) > \phi(x)$  for  $x \in \Omega$ ,  $x \neq x_0$ .

Alternatively, we could require that  $u - \phi$  has a strict local minimum at  $x_0$ . If no such test function exists, nothing is required. The lower semicontinuous functions satisfying the first half of the definition are called supersolutions, and the upper semicontinuous functions satisfying the second half are called subsolutions.

Given a point  $x \in \Omega$  we consider the class of good test functions

$$A(x) = \{\phi \in C^2 \text{ with } \nabla\phi(x) \neq 0 \text{ or } D^2\phi(x) = \mathbf{0}\}.$$

When  $\phi \in A(x)$  we can always uniquely define  $\mathbb{F}(\nabla\phi(x), D^2\phi(x))$ . When  $\nabla\phi(x) = 0$  we set

$$(10) \quad \Delta_p^N \phi(x) = \lim_{\eta \rightarrow 0} \mathbb{F}_p(\eta, \mathbf{0}) = 0.$$

**Definition 2.** A continuous function  $v$  is a viscosity solution to the equation

$$\mathbb{F}_p(\nabla v, D^2v) = g(x)$$

at  $x$ , if and only if every  $C^2$ -function  $\phi \in A(x)$  that touches  $v$  from below in  $x$  satisfies

$$\mathbb{F}_p(\nabla\phi(x), D^2\phi(x)) \leq g(x)$$

and every  $C^2$ -function  $\phi \in A(x)$  that touches  $u$  from above at  $x$  satisfies

$$\mathbb{F}_p(\nabla\phi(x), D^2\phi(x)) \geq g(x).$$

To state our third definition we need the semicontinuous extensions of  $\mathbb{F}_p$ . For a symmetric matrix  $X$  we denote by  $\lambda_{\min}(X)$  and  $\lambda_{\max}(X)$  the smallest and largest eigenvalues of  $X$  respectively. For  $p > 2$  the upper-semicontinuous extension is given by

$$\mathbb{F}_p^*(0, X) = \limsup_{\eta \rightarrow 0} \mathbb{F}_p(\eta, X) = \frac{1}{p} \text{trace}(X) + \frac{(p-2)}{p} \lambda_{\max}(X)$$

and the lower-semicontinuous extension by

$$\mathbb{F}_{*,p}(0, X) = \liminf_{\eta \rightarrow 0} \mathbb{F}_p(\eta, X) = \frac{1}{p} \text{trace}(X) + \frac{(p-2)}{p} \lambda_{\min}(X).$$

Otherwise, we define  $\mathbb{F}_p^* = \mathbb{F}_{*,p} = \mathbb{F}_p$ . For  $p < 2$  we need to exchange  $\lambda_{\min}(X)$  and  $\lambda_{\max}(X)$ . Observe that

$$-\infty < \mathbb{F}_{*,p}(0, X) \leq \mathbb{F}_p^*(0, X) < \infty$$

and that for  $\phi \in A(x)$  we have

$$\mathbb{F}_{*,p}(\nabla\phi(x), D^2\phi(x)) = \mathbb{F}_p^*(\nabla\phi(x), D^2\phi(x)).$$

**Definition 3.** A continuous function  $v$  is a viscosity solution to the equation

$$\mathbb{F}_p(\nabla v, D^2v) = g(x)$$

at  $x$ , if and only if every  $C^2$ -function  $\phi$  that touches  $v$  from below in  $x$  satisfies

$$\mathbb{F}_{*,p}(\nabla\phi(x), D^2\phi(x)) \leq g(x)$$

and every  $C^2$ -function  $\phi$  that touches  $u$  from above at  $x$  satisfies

$$\mathbb{F}_p^*(\nabla\phi(x), D^2\phi(x)) \geq g(x).$$

The above definitions are equivalent. The proof of this fact is based on the well-known fourth order perturbation argument, cf. [4], [8] or [11].

**Proposition 4.** Definitions 1, 2, and 3 are equivalent for  $1 \leq p \leq \infty$  and  $g \in C(\bar{\Omega})$ ,  $g > 0$  (or  $g < 0$ ).

*Proof.* We restrict ourselves to the case of finite  $p$ , since the case  $p = \infty$  follows by a simple modification. Clearly Definitions 1 and 3 are equivalent, and we can focus attention on showing that we can restrict the class of test functions as in Definition 2. We show directly that if Definition 3 fails, then also Definition 2 fails. To this end, we suppose that there is  $\phi \in C^2(\bar{\Omega})$  and  $x_0 \in \Omega$  such that

- i)  $u(x_0) = \phi(x_0)$ ,
- ii)  $u(x) > \phi(x)$  for  $x \in \Omega$ ,  $x \neq x_0$ ,

for which  $\nabla\phi(x_0) = 0$  and

$$(11) \quad pg(x_0) < \lambda_{\min}((p-2)D^2\phi(x_0)) + \Delta\phi(x_0).$$

We then go on showing that there exists a test function  $\phi$  with either  $\nabla\phi(x) \neq 0$  or  $\nabla\phi(x) = 0$ ,  $D^2\phi(x) = \mathbf{0}$ , for which the definition of a viscosity solution fails.

Let  $\delta > 0$  be small, and set

$$w_j(x, y) = (1 - \delta)u(x) - \left( \phi(y) - \frac{j}{4}|x - y|^4 \right)$$

and denote by  $(x_j, y_j)$  the minimum point of  $w_j$  in  $\bar{\Omega} \times \bar{\Omega}$ . Since  $x_0$  is a strict local minimum for  $u - \phi$  there exists a strict local minimum  $x_0^\delta$  for  $(1 - \delta)u - \phi$  and small enough  $\delta > 0$  such that  $x_0^\delta \rightarrow x_0$  as  $\delta \rightarrow 0$ . By first choosing a small enough  $\delta > 0$  and then large enough  $j$ , we have  $x_j, y_j \in \Omega$ , and

$$x_j, y_j \rightarrow x_0^\delta, \quad \text{as } j \rightarrow \infty.$$

We observe that

$$\phi(y) - \frac{j}{4}|x_j - y|^4,$$

has a local maximum at  $y_j$ . By (11) and continuity of

$$x \mapsto \lambda_{\min}((p-2)D^2\phi(x)) + \Delta\phi(x),$$

and  $g$ , we have

$$(12) \quad pg(y_j) < \lambda_{\min}((p-2)D^2\phi(y_j)) + \Delta\phi(y_j)$$

for small enough  $\delta > 0$  and large enough  $j$ . We denote  $\varphi = \frac{j}{4}|x_j - y_j|^4$ , and observe that  $D^2\phi(y_j) \leq D^2\varphi(y_j)$ . Thus by (12) we have

$$(13) \quad pg(y_j) < \lambda_{\min}((p-2)D^2\varphi(y_j)) + \Delta\varphi(y_j).$$

This also holds when  $p < 2$ , because

$$\begin{aligned} \lambda_{\min}((p-2)D^2\phi(y_j)) + \Delta\phi(y_j) &= (p-2)\lambda_{\max}(D^2\phi(y_j)) + \text{trace}(D^2\phi(y_j)) \\ &= (p-1)\lambda_{\max}(D^2\phi(y_j)) + \sum_{\lambda_i \neq \lambda_{\max}} \lambda_i(D^2\phi(y_j)) \\ &\leq (p-1)\lambda_{\max}(D^2\varphi(y_j)) + \sum_{\lambda_i \neq \lambda_{\max}} \lambda_i(D^2\varphi(y_j)). \end{aligned}$$

We consider the two cases: either  $x_j \neq y_j$  for all  $j$  large enough or  $x_j = y_j$  infinitely often. First, let  $y_j \neq x_j$ . We use the theorem of sums for  $w_j$ , see [5] and also [6]. It implies that there exists symmetric matrices  $X_j, Y_j$  such that  $X_j - Y_j$  is positive semidefinite and

$$\begin{aligned} \left( j|x_j - y_j|^2(x_j - y_j), Y_j \right) &\in \bar{J}^{2,+}\phi(y_j) \\ \left( j|x_j - y_j|^2(x_j - y_j), X_j \right) &\in \bar{J}^{2,-}u_\delta(x_j), \end{aligned}$$

where we denoted  $u_\delta = (1-\delta)u$ . Using this fact, inequality (12), the continuity of  $g$ , and the fact that  $g > 0$  in  $\Omega$ , we get for large enough  $j$  that

$$\begin{aligned} (1-\delta)pg(x_j) &< pg(y_j) \\ &< (p-2)\left\langle Y_j \frac{(x_j - y_j)}{|x_j - y_j|}, \frac{(x_j - y_j)}{|x_j - y_j|} \right\rangle + \text{trace}(Y_j) \\ &\leq (p-2)\left\langle X_j \frac{(x_j - y_j)}{|x_j - y_j|}, \frac{(x_j - y_j)}{|x_j - y_j|} \right\rangle + \text{trace}(X_j) \end{aligned}$$

so that the definition of the viscosity solution fails already for  $(j|x_j - y_j|^2(x_j - y_j), X_j) \in \bar{J}^{2,-}u_\delta(x_j)$  with nonvanishing  $j|x_j - y_j|^2(x_j - y_j)$ . In the case  $g < 0$  in  $\Omega$  we need to replace  $1 - \delta$  by  $1 + \delta$  throughout the argument. If  $p < 2$ , the

last inequality follows from the calculation

$$\begin{aligned}
& (p-2) \left\langle (Y_j - X_j) \frac{(x_j - y_j)}{|x_j - y_j|}, \frac{(x_j - y_j)}{|x_j - y_j|} \right\rangle + \text{trace}(Y_j - X_j) \\
& \leq (p-2)\lambda_{\min} + \sum_{i=1}^n \lambda_i \\
& = (p-1)\lambda_{\min} + \sum_{\lambda_i \neq \lambda_{\min}} \lambda_i \\
& \leq 0,
\end{aligned}$$

where  $\lambda_i$ ,  $\lambda_{\min}$ , and  $\lambda_{\max}$  denote the eigenvalues of  $Y_j - X_j$ .

Let then  $x_j = y_j$ . The fact  $D^2\varphi(y_j) = D^2\left(\frac{j}{4}|x_j - y_j|^4\right) = 0$  together with (13) shows that this case cannot happen. If  $g$  were negative instead of positive, this case would show that there exists a test function with  $\nabla\varphi(y_j) = 0$ ,  $D^2\varphi(y_j) = \mathbf{0}$  for which Definition 2 fails.  $\square$

A similar argument also provides comparison principle and uniqueness, see also Lu and Wang [17, 18]. Notice that Theorem 5 below is only stated for  $g > 0$ . In fact, for  $p = 1$  and  $g \equiv 0$ , there is a counterexample in [25], see also [13] and [14].

**Theorem 5.** *Let  $\Omega$  be a bounded domain,  $u$  lower semicontinuous and  $v$  upper semicontinuous. Suppose that  $v$  is a subsolution, and  $u$  a supersolution to (9) with  $g \in C(\bar{\Omega})$ ,  $g > 0$  and  $1 \leq p \leq \infty$ . Further, suppose that  $v \leq u$  on  $\partial\Omega$  in the sense that*

$$(14) \quad \limsup_{x \rightarrow z} v(x) \leq \liminf_{x \rightarrow z} u(x)$$

for all  $z \in \partial\Omega$ , where both sides are not simultaneously  $-\infty$  or  $\infty$ . Then

$$v \leq u \quad \text{in } \Omega.$$

*Proof.* We consider first the case  $2 \leq p < \infty$ . We argue by contradiction and assume that  $u - v$  has a strict interior minimum, that is,

$$u(x_0) - v(x_0) = \inf_{\Omega} (u - v) < \inf_{\partial\Omega} (u - v).$$

Let  $\delta \in (0, 1)$ , and set

$$w_j(x, y) = (1 - \delta)u(x) - \left(v(y) - \frac{j}{4}|x - y|^4\right)$$

and denote by  $(x_j, y_j)$  the minimum point of  $w_j$  in  $\bar{\Omega} \times \bar{\Omega}$ . Since  $x_0$  is a local minimum for  $u - v$ , there exists a strict local minimum  $x_0^\delta$  for  $(1 - \delta)u - v$  such that  $x_0^\delta \rightarrow x_0$  as  $\delta \rightarrow 0$ . Further

$$x_j, y_j \rightarrow x_0^\delta, \quad \text{as } j \rightarrow \infty$$



and  $x_j, y_j \in \Omega$  for all large  $j$ . It follows that

$$y \mapsto v(y) - \frac{j}{4} |x_j - y|^4,$$

has a local maximum at  $y_j$ , and

$$x \mapsto (1 - \delta)u(x) + \frac{j}{4} |x - y_j|^4,$$

a local minimum at  $x_j$ .

Observe that if  $y_j = x_j$ , then  $\nabla\varphi(y_j) = 0$ ,  $D^2\varphi(y_j) = \mathbf{0}$ , which immediately contradicts with the subsolution property of  $v$  since  $g > 0$ . Thus we may concentrate on the case  $x_j \neq y_j$ . Again, theorem of sums for  $w_j$  implies that there exists symmetric matrices  $X_j, Y_j$  such that  $X_j - Y_j$  is positive semidefinite and

$$\begin{aligned} \left( j |x_j - y_j|^2 (x_j - y_j), Y_j \right) &\in \bar{\mathcal{J}}^{2,+} v(y_j) \\ \left( j |x_j - y_j|^2 (x_j - y_j), X_j \right) &\in \bar{\mathcal{J}}^{2,-} u_\delta(x_j), \end{aligned}$$

where we denoted  $u_\delta = (1 - \delta)u$  so that  $u_\delta$  satisfies  $(1 - \delta)g \geq \Delta_p^N u_\delta$  in the viscosity sense i.e.

$$(1 - \delta)p g(x_j) \geq (p - 2) \left\langle X_j \frac{(x_j - y_j)}{|x_j - y_j|}, \frac{(x_j - y_j)}{|x_j - y_j|} \right\rangle + \text{trace}(X_j).$$

Using this and the corresponding inequality for  $v$ , we get for large enough  $j$  that

$$\begin{aligned} 0 &< p g(y_j) - (1 - \delta)p g(x_j) \\ &\leq (p - 2) \left\langle Y_j \frac{(x_j - y_j)}{|x_j - y_j|}, \frac{(x_j - y_j)}{|x_j - y_j|} \right\rangle + \text{trace}(Y_j) \\ &\quad - (p - 2) \left\langle X_j \frac{(x_j - y_j)}{|x_j - y_j|}, \frac{(x_j - y_j)}{|x_j - y_j|} \right\rangle - \text{trace}(X_j) \\ &= (p - 2) \left\langle (Y_j - X_j) \frac{(x_j - y_j)}{|x_j - y_j|}, \frac{(x_j - y_j)}{|x_j - y_j|} \right\rangle + \text{trace}(Y_j - X_j) \\ &\leq 0, \end{aligned}$$

because  $Y_j - X_j$  is negative semidefinite. In the first inequality we used continuity of  $g$ . This provides the desired contradiction. The cases  $1 \leq p < 2$  and  $p = \infty$  can be treated in a similar fashion as above.  $\square$

According to [12], when  $1 < p < \infty$  and  $g \equiv 0$ , it is enough to test using test functions with  $\nabla\phi(x) \neq 0$ . This definition still guarantees the uniqueness. We observe that the proof of Proposition 4 shows that in the case  $g \equiv 0$  and  $1 \leq p \leq \infty$ , Definition 3 is equivalent to a definition where we only use test functions with  $\nabla\phi(x) \neq 0$ .

**Theorem 6.** *A continuous function  $v$  is a viscosity solution to the equation*

$$\mathbb{F}_p(\nabla v, D^2 v) = 0, \quad 1 \leq p \leq \infty$$

*at  $x$ , if and only if every  $C^2$ -function  $\phi$  with  $\nabla\phi(x) \neq 0$  that touches  $v$  from below in  $x$  satisfies*

$$\mathbb{F}_p(\nabla\phi(x), D^2\phi(x)) \leq 0$$

*and every  $C^2$ -function  $\phi$ ,  $\nabla\phi(x) \neq 0$  that touches  $u$  from above at  $x$  satisfies*

$$\mathbb{F}_p(\nabla\phi(x), D^2\phi(x)) \geq 0.$$

### 3. 1-HARMONIC FUNCTIONS IN THE SENSE OF AVERAGES

Given a unit vector  $\nu \in \mathbb{R}^n$  consider the  $(n-1)$ -dimensional hyperplane

$$\pi = \nu^\perp = \{x \in \mathbb{R}^n : \langle x, \nu \rangle = 0\}.$$

For small  $\varepsilon > 0$  we denote by  $B_\varepsilon^\pi$  the  $(n-1)$ -dimensional ball in  $\pi$  centered at 0 with radius  $\varepsilon$

$$B_\varepsilon^\pi = B_\varepsilon(0) \cap \pi.$$

Let  $\Omega \subset \mathbb{R}^n$  be an open set and  $u: \Omega \rightarrow \mathbb{R}$  be a  $C^2$ -function.

Averaging the Taylor expansion

$$(15) \quad u(x+h) = u(x) + \langle \nabla u(x), h \rangle + \frac{1}{2} \langle D^2 u(x) h, h \rangle + o(|h|^2),$$

over  $B_\varepsilon^\pi$  we obtain

$$(16) \quad \int_{B_\varepsilon^\pi} u(x+h) d\mathcal{L}^{n-1}(h) = u(x) + \varepsilon^2 \cdot \frac{1}{2(n+1)} \Delta_\pi u(x) + o(\varepsilon^2),$$

where  $\Delta_\pi$  denotes the Laplace operator on the plane  $x + \pi$ . To see this, we use the orthonormal basis made up of  $\nu$  and an orthonormal basis for  $\pi$ , and observe that

$$\frac{1}{2} \int_{B_\varepsilon^\pi} \langle D^2 u(x) h, h \rangle d\mathcal{L}^{n-1}(h) = \varepsilon^2 \cdot \frac{1}{2(n+1)} \Delta_\pi u(x),$$

cf. [19]. We denote by  $D_{\nu\nu}^2 u(x) = \langle D^2 u(x) \nu, \nu \rangle$  the second derivative of  $u$  at  $x$  in the direction  $\nu$ . Note that

$$\begin{aligned} \Delta u(x) &= \text{trace}(D^2 u(x)) = \Delta_\pi u(x) + D_{\nu\nu}^2 u(x) \\ &= \Delta_\pi u(x) + \langle D^2 u(x) \nu, \nu \rangle. \end{aligned}$$

Thus we get a formula for  $\Delta_\pi$

$$(17) \quad \Delta_\pi u(x) = \Delta u(x) - \langle D^2 u(x) \nu, \nu \rangle.$$

Suppose that  $\nabla u(x) \neq 0$  and write

$$\nu = -\frac{\nabla u(x)}{|\nabla u(x)|}.$$

The vector  $\nu$  is the exterior normal to the level set

$$S = \{y \in \mathbb{R}^n : u(y) \geq u(x)\}.$$

Whenever  $\nabla u(x)$  is nonzero, the mean curvature  $H(x)$  of  $S$  is given by

$$H(x) = \frac{1}{n-1} \operatorname{div}(-\nu),$$

so that we can rewrite  $\Delta_\pi u(x)$  as

$$(18) \quad \begin{aligned} \Delta_\pi u(x) &= \Delta_1^N u(x) = |\nabla u(x)| \operatorname{div} \left( \frac{\nabla u}{|\nabla u|} \right) (x) \\ &= (n-1)H(x) |\nabla u(x)|. \end{aligned}$$

Here  $\Delta_1^N$  refers to the notation introduced in (8).

Equation (16) immediately implies a characterization of harmonic functions on the hyperplane in a sense of averages.

**Proposition 7.** *Let  $u \in C^2(\Omega)$ ,  $\nu \in \mathbb{R}^n$  be a unit vector, and  $\pi$  the  $(n-1)$ -dimensional hyperplane defined by  $\nu$ . Then*

$$\int_{B_\varepsilon^\pi} u(x+h) d\mathcal{L}^{n-1}(h) = u(x) + o(\varepsilon^2)$$

if and only if  $\Delta_\pi u(x) = 0$ .

We define unit vectors  $\nu_{\min}$  and  $\nu_{\max}$  by requiring that

$$(19) \quad \begin{aligned} u(x + \varepsilon \nu_{\min}) &= \min_{y \in \overline{B_\varepsilon}(x)} u(y) \\ u(x + \varepsilon \nu_{\max}) &= \max_{y \in \overline{B_\varepsilon}(x)} u(y). \end{aligned}$$

Observe that whenever  $\nabla u(x) \neq 0$ , then  $\nu_{\min}$  and  $\nu_{\max}$  converge to the uniquely defined directions

$$-\frac{\nabla u(x)}{|\nabla u(x)|} \quad \text{and} \quad \frac{\nabla u(x)}{|\nabla u(x)|}$$

respectively even if those vectors themselves may not be unique.

**Definition 8.** *A continuous function  $u$  is 1-harmonic in the sense of averages if*

$$(20) \quad u(x) = \int_{B_\varepsilon^{\pi\nu}} u(x+h) d\mathcal{L}^{n-1}(h) + o(\varepsilon^2) \quad \text{as } \varepsilon \rightarrow 0,$$

in the viscosity sense *i.e.*

(1) *if for every  $\phi \in C^2$  such that  $\phi$  touches  $u$  at  $x \in \Omega$  from below, we have*

$$\phi(x) \geq \int_{B_\varepsilon^{\pi\nu_{\max}}} \phi(x+h) d\mathcal{L}^{n-1}(h) + o(\varepsilon^2),$$

*for any  $\nu_{\max}$  in (19) as  $\varepsilon \rightarrow 0$ .*

(2) if for every  $\phi \in C^2$  such that  $\phi$  touches  $u$  at  $x \in \Omega$  from above, we have

$$\phi(x) \leq \int_{B_\varepsilon^{\pi\nu_{\min}}} \phi(x+h) d\mathcal{L}^{n-1}(h) + o(\varepsilon^2),$$

for any  $\nu_{\min}$  in (19) as  $\varepsilon \rightarrow 0$ .

**Theorem 9.** A continuous function  $u$  in a domain  $\Omega \subset \mathbb{R}^n$  is 1-harmonic in the sense of averages if and only if

$$\Delta_1^N u = 0$$

in the viscosity sense.

*Proof.* For a smooth  $\phi$

$$(21) \quad \int_{B_\varepsilon^{\pi\nu}} \phi(x+h) d\mathcal{L}^{n-1}(h) = \phi(x) + \frac{\varepsilon^2}{2(n+1)} \Delta_{\pi\nu} \phi(x) + o(\varepsilon^2),$$

holds for any  $\nu \neq 0$ . If  $\nabla\phi \neq 0$ , then

$$(22) \quad \nu_{\min} \rightarrow -\nabla\phi / |\nabla\phi| = \nu \quad \text{as } \varepsilon \rightarrow 0.$$

We choose  $\phi \in C^2$ ,  $\nabla\phi \neq 0$  such that  $\phi$  touches  $u$  at  $x \in \Omega$  from below, and suppose that  $u$  is a viscosity solution to  $\Delta u - \Delta_\infty^N u = \Delta_{\pi\nu} u = 0$ . In particular,  $\Delta_{\pi\nu} \phi(x) \leq 0$ . This, (21), (22), and the continuity of the second derivatives imply

$$\int_{B_\varepsilon^{\pi\nu_{\min}}} \phi(x+h) d\mathcal{L}^{n-1}(h) \leq \phi(x) + o(\varepsilon^2).$$

The second half of the definition of a viscosity solution follows similarly.

Suppose then that  $u$  is 1-harmonic in the sense of averages, i.e. for the above  $\phi$  we have

$$\phi(x) \geq \int_{B_\varepsilon^{\pi\nu_{\min}}} \phi(x+h) d\mathcal{L}^{n-1}(h) + o(\varepsilon^2).$$

Combining this together with (21) we obtain

$$0 \geq \frac{\varepsilon^2}{2(n+1)} \Delta_{\pi\nu_{\min}} \phi(x) + o(\varepsilon^2).$$

Dividing this by  $\varepsilon^2$ , passing to a limit with  $\varepsilon$ , and using (22), we see that  $u$  satisfies the condition for the viscosity supersolution with this  $\phi$ . The proof for the second half is analogous.

We are left with the case  $\nabla\phi(x) = 0$ . Suppose that  $\phi \in C^2$  such that  $\phi$  touches  $u$  at  $x \in \Omega$  from below with  $\nabla\phi(x) = 0$ , and suppose that  $u$  is a viscosity solution so that

$$-\lambda_{\max}(D^2\phi(x)) + \Delta\phi(x) \leq 0.$$

Observing that now

$$(23) \quad \langle D^2 \phi \nu_{\max}, \nu_{\max} \rangle \rightarrow \lambda_{\max}(D^2 \phi(x))$$

as  $\varepsilon \rightarrow 0$ , and combining this with (21), we see that  $u$  is 1-harmonic in the sense of averages.

Suppose then that  $u$  is 1-harmonic in the sense of averages, i.e. for the above  $\phi$  we have

$$\phi(x) \geq \int_{B_\varepsilon^{\pi \nu_{\max}}} \phi(x+h) d\mathcal{L}^{n-1}(h) + o(\varepsilon^2).$$

Combining this together with (21) and (23), dividing by  $\varepsilon^2$  and passing to a limit with  $\varepsilon$ , we see that  $u$  satisfies the first half of a definition of a viscosity solution. The second half is again analogous.  $\square$

#### 4. $p$ -HARMONIC FUNCTIONS IN THE SENSE OF AVERAGES

We start with a formal calculation assuming that  $u$  is smooth and  $\nabla u \neq 0$ . The gradient direction is almost the maximizing direction for a smooth function whenever the gradient does not vanish. We insert  $h = \pm \nabla u / |\nabla u|$  in (15) and sum up the two resulting expansions to get rid of the first order terms

$$(24) \quad \begin{aligned} & u(x) - \frac{1}{2} \left\{ \max_{\bar{B}_\varepsilon(x)} u + \min_{\bar{B}_\varepsilon(x)} u \right\} \\ & \approx u(x) - \frac{1}{2} \left\{ u \left( x + \varepsilon \frac{\nabla u(x)}{|\nabla u(x)|} \right) + u \left( x - \varepsilon \frac{\nabla u(x)}{|\nabla u(x)|} \right) \right\} \\ & = -\frac{\varepsilon^2}{2} \Delta_\infty^N u(x) + o(\varepsilon^2). \end{aligned}$$

Next we multiply (24) and (16) by the constants  $\alpha = \frac{p-1}{p+n}$  and  $\beta = \frac{n+1}{p+n}$  satisfying  $\alpha + \beta = 1$  and add up the formulas so that we have the operator in (6) on the right hand side. We get

$$\begin{aligned} & \frac{n+1}{p+n} \int_{B_\varepsilon^{\pi \nu_{\min}}} u(x+h) d\mathcal{L}^{n-1}(h) + \left( \frac{p-1}{p+n} \right) \frac{\max_{\bar{B}_\varepsilon(x)} u + \min_{\bar{B}_\varepsilon(x)} u}{2} \\ & \approx u(x) + \frac{\varepsilon^2}{2(p+n)} ((p-1) \Delta_\infty^N u + \Delta_1^N u(x)) + o(\varepsilon^2). \end{aligned}$$

This motivates the following definition which we only formulate in the case  $1 \leq p \leq 2$ . In the case  $p > 2$  the definition is almost identical, except that  $\nu_{\min}$  and  $\nu_{\max}$  should be interchanged. This only plays a role when  $\nabla \phi$  vanishes. In that case

$$\langle D^2 \phi \nu_{\max}, \nu_{\max} \rangle \rightarrow \lambda_{\max}(D^2 \phi(x)) \quad \text{and} \quad \langle D^2 \phi \nu_{\min}, \nu_{\min} \rangle \rightarrow \lambda_{\min}(D^2 \phi(x))$$

see also (23).

**Definition 10.** A continuous function  $u$  is  $p$ -harmonic,  $1 \leq p \leq 2$ , in the sense of averages if it satisfies

$$(25) \quad u(x) = \frac{n+1}{p+n} \int_{B_\varepsilon^{\pi\nu}} u(x+h) d\mathcal{L}^{n-1}(h) + \left(\frac{p-1}{p+n}\right) \frac{\max_{\overline{B_\varepsilon(x)}} u + \min_{\overline{B_\varepsilon(x)}} u}{2} + o(\varepsilon^2),$$

as  $\varepsilon \rightarrow 0$  in the viscosity sense, that is,

1. for every  $\phi \in C^2$  such that  $\phi$  touches  $u$  at  $x \in \Omega$  from below, we have

$$\phi(x) \geq \frac{n+1}{p+n} \int_{B_\varepsilon^{\pi\nu_{\max}}} \phi(x+h) d\mathcal{L}^{n-1}(h) + \left(\frac{p-1}{p+n}\right) \frac{\max_{\overline{B_\varepsilon(x)}} \phi + \min_{\overline{B_\varepsilon(x)}} \phi}{2} + o(\varepsilon^2),$$

for any  $\nu_{\max}$  in (19) as  $\varepsilon \rightarrow 0$ ,

2. for every  $\phi \in C^2$  such that  $\phi$  touches  $u$  at  $x \in \Omega$  from above, we have

$$\phi(x) \leq \frac{n+1}{p+n} \int_{B_\varepsilon^{\pi\nu_{\min}}} \phi(x+h) d\mathcal{L}^{n-1}(h) + \left(\frac{p-1}{p+n}\right) \frac{\max_{\overline{B_\varepsilon(x)}} \phi + \min_{\overline{B_\varepsilon(x)}} \phi}{2} + o(\varepsilon^2),$$

for any  $\nu_{\min}$  in (19) as  $\varepsilon \rightarrow 0$ .

Unfortunately, using test functions in the above definition instead of  $u$  itself seems to be necessary to obtain the next theorem as indicated by the known counterexample in the case  $p = \infty$  (see [19] for an example).

For  $p = 1$ , Theorem 11 follows from Theorem 9, for  $p = \infty$ , the proof follows from (5), cf. Theorem 2 in [19], and for  $p \in (1, \infty)$  it is analogous to the proof of Theorem 13 below, so we omit it here.

**Theorem 11.** Let  $1 \leq p \leq \infty$ . A continuous function  $u$  in a domain  $\Omega \subset \mathbb{R}^n$  is  $p$ -harmonic in the sense of averages according to Definition 10 if and only if

$$\Delta_p^N u(x) = 0$$

in the viscosity sense.

## 5. MEAN VALUE FORMULA FOR A TUG-OF-WAR GAME

We already defined  $p$ -harmonic functions in the sense of averages in the previous section. In this section, we derive another mean value formula. It appears to be more complicated, but in the context of the tug-of-war game similar to that in [22] it turns out to be quite natural. Below

$$\alpha = \frac{p-1}{p+n} \quad \beta = \frac{n+1}{p+n},$$

and  $B_\varepsilon^{\pi\nu}$  is the  $(n-1)$ -dimensional ball of radius  $\varepsilon$  centered at zero in the hyperplane  $\pi_\nu$ .

**Definition 12.** A continuous function  $u$  satisfies

$$(26) \quad u(x) = \frac{1}{2} \sup_{0 < |\nu| \leq 1} \left\{ \alpha u(x + \nu\varepsilon) + \beta \int_{B_\varepsilon^{\pi\nu}} u(x+h) d\mathcal{L}^{n-1}(h) \right\} \\ + \frac{1}{2} \inf_{0 < |\nu| \leq 1} \left\{ \alpha u(x + \nu\varepsilon) + \beta \int_{B_\varepsilon^{\pi\nu}} u(x+h) d\mathcal{L}^{n-1}(h) \right\} + o(\varepsilon^2),$$

as  $\varepsilon \rightarrow 0$  in the sense of averages if

(1) for every  $\phi \in C^2$  such that  $\phi$  touches  $u$  at  $x \in \Omega$  from below with  $\nabla\phi(x) \neq 0$ , we have

$$\phi(x) \geq \frac{1}{2} \sup_{0 < |\nu| \leq 1} \left\{ \alpha\phi(x + \nu\varepsilon) + \beta \int_{B_\varepsilon^{\pi\nu}} \phi(x+h) d\mathcal{L}^{n-1}(h) \right\} \\ + \frac{1}{2} \inf_{0 < |\nu| \leq 1} \left\{ \alpha\phi(x + \nu\varepsilon) + \beta \int_{B_\varepsilon^{\pi\nu}} \phi(x+h) d\mathcal{L}^{n-1}(h) \right\} + o(\varepsilon^2).$$

(2) for every  $\phi \in C^2$  such that  $\phi$  touches  $u$  at  $x \in \Omega$  from above with  $\nabla\phi(x) \neq 0$ , we have

$$\phi(x) \leq \frac{1}{2} \sup_{0 < |\nu| \leq 1} \left\{ \alpha\phi(x + \nu\varepsilon) + \beta \int_{B_\varepsilon^{\pi\nu}} \phi(x+h) d\mathcal{L}^{n-1}(h) \right\} \\ + \frac{1}{2} \inf_{0 < |\nu| \leq 1} \left\{ \alpha\phi(x + \nu\varepsilon) + \beta \int_{B_\varepsilon^{\pi\nu}} \phi(x+h) d\mathcal{L}^{n-1}(h) \right\} + o(\varepsilon^2).$$

The case  $p = \infty$  is already considered in [19], and thus we concentrate on the case  $1 < p < \infty$ .

**Theorem 13.** Let  $1 < p < \infty$ . A continuous function  $u$  in a domain  $\Omega \subset \mathbb{R}^n$  satisfies Definition 12 if and only if

$$\Delta_p^N u(x) = 0$$

in the viscosity sense.

*Proof.* First we recall a calculation from [19] leading to an asymptotic expansion involving the infinity Laplacian. Choose a point  $x \in \Omega$  and a  $C^2$ -function  $\phi$  defined in a neighborhood of  $x$ . Let  $\nu_{\min}$  be a vector giving

$$\inf_{0 < |\nu| \leq 1} \left\{ \alpha\phi(x + \nu\varepsilon) + \beta \int_{B_\varepsilon^{\pi\nu}} \phi(x+h) d\mathcal{L}^{n-1}(h) \right\}.$$

Later we check that the infimum is not obtained when  $\nu \rightarrow 0$  so that  $\nu_{\min}$  really exists. Similarly, let  $\nu_{\max}$  be a vector giving

$$\sup_{0 < |\nu| \leq 1} \left\{ \alpha\phi(x + \nu\varepsilon) + \beta \int_{B_\varepsilon^{\pi\nu}} \phi(x+h) d\mathcal{L}^{n-1}(h) \right\}.$$

Consider the Taylor expansion of the second order of  $\phi$

$$\phi(x + \nu) = \phi(x) + \nabla\phi(x) \cdot \nu + \frac{1}{2}\langle D^2\phi(x)\nu, \nu \rangle + o(|\nu|^2)$$

as  $|\nu| \rightarrow 0$ . Evaluating this Taylor expansion of  $\phi$  at the point  $x$  with  $\nu = \nu_{\min}\varepsilon$ , and  $\nu = -\nu_{\min}\varepsilon$  we get

$$(27) \quad \phi(x + \nu_{\min}\varepsilon) = \phi(x) + \nabla\phi(x) \cdot \nu_{\min}\varepsilon + \frac{1}{2}\langle D^2\phi(x)\nu_{\min}\varepsilon, \nu_{\min}\varepsilon \rangle + o(\varepsilon^2),$$

$$(28) \quad \phi(x - \nu_{\min}\varepsilon) = \phi(x) - \nabla\phi(x) \cdot \nu_{\min}\varepsilon + \frac{1}{2}\langle D^2\phi(x)\nu_{\min}\varepsilon, \nu_{\min}\varepsilon \rangle + o(\varepsilon^2)$$

as  $\varepsilon \rightarrow 0$ . Adding the expressions, we obtain

$$(29) \quad \phi(x - \nu_{\min}\varepsilon) + \phi(x + \nu_{\min}\varepsilon) - 2\phi(x) = \varepsilon^2\langle D^2\phi(x)\nu_{\min}, \nu_{\min} \rangle + o(\varepsilon^2).$$

We also have

$$(30) \quad \int_{B_\varepsilon^{\pi\nu}} \phi(x + h) d\mathcal{L}^{n-1}(h) = \phi(x) + \frac{\varepsilon^2}{2(n+1)}\Delta_{\pi\nu}\phi(x) + o(\varepsilon^2),$$

for any nonzero  $\nu$ . We choose  $\nu = \nu_{\min}$  and  $\nu = -\nu_{\min}$  in (30), multiply by  $\alpha/2$  and sum up the results with (29) multiplied by  $\beta/2$ . Then we use the definition of  $\nu_{\max}$  and estimate  $\phi(x - \nu_{\min}\varepsilon) + \int_{B_\varepsilon^{\pi-\nu_{\min}}} \phi(x + h) d\mathcal{L}^{n-1}(h)$  from above, as well as observe that  $\Delta_{\pi\nu_{\min}} = \Delta_{\pi-\nu_{\min}}$ , and obtain

$$(31) \quad \begin{aligned} & \frac{1}{2} \sup_{0 < |\nu| \leq 1} \left\{ \alpha\phi(x + \nu\varepsilon) + \beta \int_{B_\varepsilon^{\pi\nu}} \phi(x + h) d\mathcal{L}^{n-1}(h) \right\} \\ & + \frac{1}{2} \inf_{0 < |\nu| \leq 1} \left\{ \alpha\phi(x + \nu\varepsilon) + \beta \int_{B_\varepsilon^{\pi\nu}} \phi(x + h) d\mathcal{L}^{n-1}(h) \right\} \\ & \geq \phi(x) \\ & + \frac{\beta\varepsilon^2}{2(n+1)} \left( (p-1)\langle D^2\phi(x)\nu_{\min}, \nu_{\min} \rangle + \Delta_{\pi\nu_{\min}}\phi(x) \right) \\ & + o(\varepsilon^2), \end{aligned}$$

which holds for any smooth function for which  $\nu_{\min} \neq 0$ .

Suppose then that  $\nabla\phi(x) \neq 0$ . By considering the lowest order terms in the sum of (27) and (30)

$$\begin{aligned} & \alpha\phi(x + \nu_{\min}\varepsilon) + \beta \int_{B_\varepsilon^{\pi\nu_{\min}}} \phi(x + h) d\mathcal{L}^{n-1}(h) = \phi(x) + \alpha\nabla\phi(x) \cdot \nu_{\min}\varepsilon \\ & + \alpha\frac{1}{2}\langle D^2\phi(x)\nu_{\min}\varepsilon, \nu_{\min}\varepsilon \rangle + \beta\frac{\varepsilon^2}{2(n+1)}\Delta_{\pi\nu_{\min}}\phi(x) + o(\varepsilon^2), \end{aligned}$$

we see that

$$(32) \quad \nu_{\min} \rightarrow -\nabla\phi / |\nabla\phi|$$

as  $\varepsilon \rightarrow 0$  because  $p \neq 1$  i.e.  $\alpha > 0$ .



Suppose that function  $u$  satisfies Definition 12. Consider a smooth  $\phi$ ,  $\nabla\phi \neq 0$  which touches  $u$  at  $x$  from below. Combining Definition 12 together with (31), we obtain

$$0 \geq \frac{\beta\varepsilon^2}{2(n+1)} \left( (p-1)\langle D^2\phi(x)\nu_{\min}, \nu_{\min} \rangle + \Delta_{\pi_{\nu_{\min}}}\phi(x) \right) + o(\varepsilon^2).$$

Dividing this by  $\varepsilon^2$  and recalling (32), and passing to a limit, we obtain the condition in the definition for the viscosity supersolution because with  $\nu = \nabla\phi/|\nabla\phi|$  we have  $\Delta_1^N\phi(x) = \Delta_{\pi_\nu}\phi = \Delta\phi - \Delta_\infty^N\phi$ .

To prove the reverse implication, assume that  $u$  is a viscosity solution. In particular  $u$  is a subsolution. Let  $\phi$  be a smooth test function touching  $u$  at  $x \in \Omega$  from above. If  $\nabla\phi(x) \neq 0$  and we set  $\nu = \nabla\phi/|\nabla\phi|$ , it follows that

$$(33) \quad \begin{aligned} 0 &\leq (p-2)\Delta_\infty^N\phi(x) + \Delta\phi(x) = (p-1)\Delta_\infty^N\phi(x) + (\Delta\phi(x) - \Delta_\infty^N\phi(x)) \\ &= (p-1)\Delta_\infty^N\phi(x) + \Delta_{\pi_\nu}\phi(x). \end{aligned}$$

This together with (31) and (32) shows that  $\phi$  satisfies the second half in Definition 12. The other cases are similar. According to Theorem 6, we only need to test with the test functions with nonvanishing gradients, and thus the proof is complete.  $\square$

## 6. TUG-OF-WAR WITH NOISE

The asymptotic expansion in the previous section is related to the tug-of-war game with noise which is quite similar to that in [22].

Fix  $\varepsilon > 0$  and consider the following two-player zero-sum-game. At the beginning, a token is placed at a point  $x_0 \in \Omega$ . First players fix their possible moves  $\nu_I$  and  $\nu_{II}$  with  $|\nu_I|, |\nu_{II}| \leq \varepsilon$ , and toss a fair coin. If Player I wins the toss, then she tosses a biased coin. If she gets heads (with probability  $\alpha > 0$ ), the token is placed at  $x_1 = x_0 + \nu_I$ . If she gets tails (with probability  $\beta > 0$ ), then the token is placed at a random point in  $x_1 \in B_\varepsilon^{\pi_{\nu_I}}(x_0)$ . Similarly if Player II wins the toss, then he tosses a biased coin. If he gets heads (with probability  $\alpha$ ), the token is placed at  $x_1 = x_0 + \nu_{II}$ . If he gets tails (with probability  $\beta$ ), then the token is placed at a random point in  $x_1 \in B_\varepsilon^{\pi_{\nu_{II}}}(x_0)$ . The game is played until the token hits

$$\Gamma_\varepsilon = \{x \in \mathbb{R}^n \setminus \Omega : \text{dist}(x, \partial\Omega) \leq \varepsilon\}.$$

This procedure yields a possibly infinite sequence of game states  $x_0, x_1, \dots$  where every  $x_k$  is a random variable. We denote by  $x_\tau \in \Gamma_\varepsilon$  the first point in  $\Gamma_\varepsilon$  in the sequence, where  $\tau$  is the hitting time. The payoff is  $F(x_\tau)$ , where  $F: \Gamma_\varepsilon \rightarrow \mathbb{R}$  is a given measurable *payoff function*. Player I earns  $F(x_\tau)$  while Player II earns  $-F(x_\tau)$ .

A history of a game up to step  $k$  is a vector of the first  $k+1$  game states and  $k$  steps, for example,  $(x_0, \nu_1, x_1, \dots, \nu_k, x_k)$ . A strategy  $S_I$  for Player I is

a Borel function defined on the space of all histories that gives the next step for Player I

$$\nu_{k+1}^I, \quad |\nu_{k+1}^I| \leq \varepsilon$$

given a history  $h$  if Player I wins the toss. Similarly Player II plays according to a strategy  $S_{\text{II}}$

Using the Kolmogorov construction the fixed starting point  $x_0$  and the strategies  $S_{\text{I}}$  and  $S_{\text{II}}$  determine a unique probability measure  $\mathbb{P}_{S_{\text{I}}, S_{\text{II}}}^{x_0}$ .

The expected payoff, when starting from  $x_0$  and using the strategies  $S_{\text{I}}, S_{\text{II}}$ , is

$$(34) \quad \mathbb{E}_{S_{\text{I}}, S_{\text{II}}}^{x_0}[F(x_\tau)] = \int_{H^\infty} F(x_\tau(\omega)) d\mathbb{P}_{S_{\text{I}}, S_{\text{II}}}^{x_0}(\omega)$$

where we integrate over all histories  $H^\infty$ .

The *value of the game for Player I* is given by

$$u_{\text{I}}^\varepsilon(x_0) = \sup_{S_{\text{I}}} \inf_{S_{\text{II}}} \mathbb{E}_{S_{\text{I}}, S_{\text{II}}}^{x_0}[F(x_\tau)]$$

while the *value of the game for Player II* is given by

$$u_{\text{II}}^\varepsilon(x_0) = \inf_{S_{\text{II}}} \sup_{S_{\text{I}}} \mathbb{E}_{S_{\text{I}}, S_{\text{II}}}^{x_0}[F(x_\tau)].$$

If the chosen strategies result in a game that does not end almost surely we set the expected pay-off for Player I to be  $-\infty$  and for Player II to be  $+\infty$ . The values  $u_{\text{I}}^\varepsilon(x_0)$  and  $u_{\text{II}}^\varepsilon(x_0)$  are intuitively the best expected outcomes each player can guarantee when the game starts at  $x_0$ .

We start by the statement of the *Dynamic Programming Principle* (DPP) applied to our game.

**Lemma 14** (DPP). *For  $\alpha > 0$  and  $\beta > 0$  (which corresponds to  $1 < p < \infty$ ) the value function for Player I satisfies*

$$(35) \quad \begin{aligned} u_{\text{I}}^\varepsilon(x) = & \frac{1}{2} \sup_{0 < |\nu| \leq 1} \left\{ \alpha u_{\text{I}}^\varepsilon(x + \nu\varepsilon) + \beta \int_{B_\varepsilon^{\pi\nu}} u_{\text{I}}^\varepsilon(x + h) d\mathcal{L}^{n-1}(h) \right\} \\ & + \frac{1}{2} \inf_{0 < |\nu| \leq 1} \left\{ \alpha u_{\text{I}}^\varepsilon(x + \nu\varepsilon) + \beta \int_{B_\varepsilon^{\pi\nu}} u_{\text{I}}^\varepsilon(x + h) d\mathcal{L}^{n-1}(h) \right\} \end{aligned}$$

for each  $x_0 \in \Omega$  and

$$u_{\text{I}}^\varepsilon(x_0) = F(x_0), \quad \text{for } x_0 \in \Gamma_\varepsilon.$$

The value function for Player II,  $u_{\text{II}}^\varepsilon$ , satisfies the same equation.

An intuitive explanation for DPP can be obtained by considering the different outcomes of a single game round with the corresponding probabilities.

It turns out that the values of the game satisfy a *comparison principle*, the values are *unique*,  $u_{\text{I}} = u_{\text{II}}$  with fixed boundary values and any function satisfying (35) is a game value. In a smooth domain with regular boundary

data, the values converge to a unique  $p$ -harmonic function as the step size tends to zero. The proofs are similar to those in [21] and [22].

**Acknowledgement:** This collaboration started in the fall of 2008 during a visit of BK to the University of Pittsburgh, where he enjoyed activities in the Analysis seminar. MP would like to thank for the hospitality during his visits to Cologne and Pittsburgh. JM research is partially supported by NSF award DMS-1001179, and MP by Academy of Finland. The authors would like to thank Petri Juutinen for useful comments.

## REFERENCES

- [1] W. Blaschke, Ein Mittelwertsatz und eine kennzeichnende Eigenschaft des logarithmischen Potentials, *Leipz. Ber.* 68:3–7, 1916.
- [2] A. Bonfiglioli and E. Lanconelli, Subharmonic Functions in sub-Riemannian Settings, *J. Europ. Math. Soc.*, to appear.
- [3] A. Bonfiglioli, E. Lanconelli, and F. Uguzzoni. *Stratified Lie groups and potential theory for their sub-Laplacians*, Springer, 2007.
- [4] Y. Chen, Y. Giga, and S. Goto, Uniqueness and existence of viscosity solutions of generalized mean curvature flow equations. *J. Differential Geom.*, 33:749–786, 1991.
- [5] M. Crandall, and H. Ishii. The maximum principle for semicontinuous functions. *Differential Integral Equations.*, 3(6):1001–1014, 1990.
- [6] M. Crandall, H. Ishii, and P.-L. Lions. User’s guide to viscosity solutions of second order partial differential equations. *Bull. Amer. Math. Soc.*, 27:1–67, 1992.
- [7] K. Does, An evolution equation involving the normalized  $p$ -Laplacian *Comm. Pure Appl. Anal.* 10:361–396, 2011. Dissertation under the same title, University of Cologne, 2009.
- [8] L. C. Evans and J. Spruck, *Motion of level sets by mean curvature I*, *J. Differential Geom.*, 33:635–681, 1991.
- [9] Y. Giga. *Surface evolution equations, A level set approach*. Vol. 99 of *Monographs in Mathematics*. Birkhäuser Verlag, Basel, 2006.
- [10] P. Juutinen.  $p$ -harmonic approximation of functions of least gradient. *Indiana Univ. Math. J.*, 54:1015–1029, 2005.
- [11] P. Juutinen and B. Kawohl, On the evolution governed by the infinity Laplacian, *Math. Annalen*, 335:819–851, 2006.
- [12] P. Juutinen, P. Lindqvist, and J. J. Manfredi. On the equivalence of viscosity solutions and weak solutions for a quasi-linear equation. *SIAM J. Math. Anal.*, 33(3):699–717, 2001.
- [13] B. Kawohl, Variations on the  $p$ -Laplacian, in: *Nonlinear Elliptic Partial Differential Equations*, Contemporary Mathematics, 540:35–46, 2011.
- [14] B. Kawohl and N. Kutev, Strong maximum principle for semicontinuous viscosity solutions of nonlinear partial differential equations, *Arch. Math.*, 70:470–478, 1998.
- [15] E. Le Gruyer. On absolutely minimizing Lipschitz extensions and PDE  $\Delta_\infty(u) = 0$ . *NoDEA Nonlinear Differential Equations Appl.*, 14(1-2):29–55, 2007.
- [16] E. Le Gruyer and J. C. Archer. Harmonious extensions. *SIAM J. Math. Anal.*, 29(1):279–292, 1998.
- [17] G. Lu and P. Wang. A PDE perspective of the normalized infinity Laplacian. *Comm. Partial Differential Equations.*, 33(10):1788–1817, 2008.
- [18] G. Lu and P. Wang. A uniqueness theorem for degenerate elliptic equations. *Lecture Notes of Seminario Interdisciplinare di Matematica, Conference on Geometric Methods*

- in PDE's (Bologna), On the Occasion of 65th Birthday of Ermanno Lanconelli. Edited by G. Citti, A. Montanari, A. Pascucci, S. Polidoro, 7:207–222, 2008.*
- [19] J. J. Manfredi, M. Parviainen, and J.D. Rossi. An asymptotic mean value characterization for  $p$ -harmonic functions. *Proc. Amer. Math. Soc.*, 258:713–728, 2010.
  - [20] J. J. Manfredi, M. Parviainen, and J.D. Rossi. An asymptotic mean value characterization for a class of nonlinear parabolic equations related to tug-of-war games. *SIAM J. Math. Anal.*, 42(5):2058–2081, 2010.
  - [21] J. J. Manfredi, M. Parviainen, and J.D. Rossi. On the definition and properties of  $p$ -harmonious functions. *Ann. Sc. Norm. Super. Pisa Cl. Sci.*, to appear
  - [22] Y. Peres and S. Sheffield. Tug-of-war with noise: a game-theoretic view of the  $p$ -Laplacian. *Duke Math. J.*, 145(1):91–120, 2008.
  - [23] Y. Peres, O. Schramm, S. Sheffield, and D. B. Wilson. Tug-of-war and the infinity Laplacian. *J. Amer. Math. Soc.*, 22(1):167–210, 2009.
  - [24] I. Privaloff, Sur les fonctions harmoniques *Rec. Math. Moscou (Mat. Sbornik)*, 32, 464–471, 1825 .
  - [25] P. Sternberg and W. P. Ziemer. Generalized motion by curvature with a Dirichlet condition. *J. Differential Equations*, 114(2):580–600, 1994.

(BK) MATHEMATICAL INSTITUTE, UNIVERSITY OF COLOGNE, 50923 COLOGNE, GERMANY, KAWOHL@MATH.UNI-KOELN.DE

(JM) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PITTSBURGH, PITTSBURGH, PA 15260, USA, MANFREDI@PITT.EDU

(MP) DEPARTMENT OF MATHEMATICS AND STATISTICS, P.O. Box 35, 40014 UNIVERSITY OF JYVÄSKYLÄ, FINLAND, MIKKO.J.PARVIAINEN@JYU.FI