

# *BLD*-MAPPINGS IN $W^{2,2}$ ARE LOCALLY INVERTIBLE

JUHA HEINONEN AND TERO KILPELÄINEN

February 18, 1999

**ABSTRACT.** We prove that mappings of bounded length distortion are local homeomorphisms if they have  $L^2$ -integrable weak second derivatives.

## 1. Introduction

In this note, we establish the following theorem:

**1.1. Theorem.** *Every  $BLD$ -mapping that belongs to the Sobolev space  $W^{2,2}$  is a local homeomorphism.*

Mappings of bounded length distortion, abbreviated *BLD*-mappings, were introduced and studied by Martio and Väisälä in [MV]: a mapping  $f$  from an open subset  $\Omega$  of  $\mathbf{R}^n$ ,  $n \geq 2$ , into  $\mathbf{R}^n$  is a *BLD*-mapping if

$$(1.2) \quad f \in L^{1,\infty}(\Omega)$$

and

$$(1.3) \quad \det df(x) \geq c > 0$$

for some constant  $c$  and for almost every  $x$  in  $\Omega$ . Here  $L^{1,\infty}(\Omega)$  is the Sobolev space of (continuous) functions with essentially bounded first distributional derivatives; thus (1.2) is equivalent to the requirement that  $f$  is locally uniformly Lipschitz: there is a constant  $L \geq 1$  such that

$$(1.4) \quad |f(x) - f(y)| \leq L|x - y|$$

---

The research of the first author is financed by the U.S. National Science Foundation (Grant DMS 96/22844). The research of the second author is financed by the Academy of Finland (Project #8597).

whenever  $x$  and  $y$  lie in a ball contained in  $\Omega$ . Lipschitz functions are almost everywhere differentiable, and in (1.3)  $df(x)$  denotes the total derivative of  $f$ .

*BLD*-mappings form an interesting subclass of general *quasiregular mappings* [Re], [Ri], and they can be characterized by a quasipreserving property of lengths of paths [MV]. (See also [HKM, Chapter 14] and [HR].)

Theorem 1.1 was informally conjectured by Dennis Sullivan in a discussion with the first author. The conjecture was based on the philosophy of [S], and corroborated by the fact that the “winding map”  $(r, \theta, w) \mapsto (r, 2\theta, w)$  in cylindrical coordinates belongs (locally) to  $W^{2,p}$  for each  $p < 2$  and fails to be a local homeomorphism on the  $(n-2)$ -dimensional (linear) subspace  $\{r=0\}$  of  $\mathbf{R}^n$ ,  $n \geq 2$ .

We recall another conjecture (due to Olli Martio) which predicts the extremality of the winding map: the so-called inner dilatation  $K_I$  of the winding map is 2 and, conjecturally, in dimensions  $n \geq 3$ , every nonconstant quasiregular map with inner dilatation strictly less than 2 is a local homeomorphism. (See [Ri, I.3.1].) In Section 3 below we shall show that the conjecture is true for quasiregular maps whose dilatation tensor belongs to the Sobolev space  $W^{1,2}$ .

## 2. Proof of Theorem 1.1

Let  $f: \Omega \rightarrow \mathbf{R}^n$ ,  $n \geq 2$ , be a mapping that satisfies (1.2) and (1.3), and assume that  $f$  has second distributional derivatives in  $L^2(\Omega)$ . By a theorem of Reshetnyak [Re, Thm. II.6.3],  $f$  is an open mapping with discrete fibers. The branch set  $B_f$  is the closed set in  $\Omega$ , where  $f$  does not define a local homeomorphism.

We assume that  $B_f \neq \emptyset$ , and then show that this leads to a contradiction. By the general theory of discrete and open mappings, the Hausdorff  $(n-2)$ -measure of  $f(B_f)$  is positive [Ri, III.5.3]; therefore, because  $f$  is locally Lipschitz, we have that

$$(2.1) \quad \mathcal{H}_{n-2}(B_f) > 0$$

as well, where  $\mathcal{H}_{n-2}$  denotes the Hausdorff  $(n-2)$ -measure in  $\mathbf{R}^n$ . (Note that (2.1) is unknown for general quasiregular mappings with nonempty branch set  $B_f$  in dimensions  $n \geq 4$ . See [Ri, III. 5.4.2] and Section 3 below.)

The key point in our argument is the following fact: there is an exceptional set  $E$  of Hausdorff  $(n-2)$ -measure zero in  $\Omega$  such that

$$(2.2) \quad \lim_{r \rightarrow 0} \int_{B(x,r)} |df(y) - (df)_{x,r}| dy = 0$$

for  $x \in \Omega \setminus E$ , where the barred integral sign denotes the integral average,  $B(x, r)$  is an open  $n$ -ball with center  $x$  and radius  $r > 0$ , and

$$(df)_{x,r} = \int_{B(x,r)} df(y) dy.$$

This fact follows from the Poincaré inequality

$$\left( \int_{B(x,r)} |df(y) - (df)_{x,r}| dy \right)^2 \leq c(n)r^2 \int_{B(x,r)} |d^2f(y)|^2 dy,$$

and the following well-known application of the Lebesgue differentiation theorem and basic covering arguments: if  $u \in L^1(\Omega)$ , then

$$\mathcal{H}_{n-2} \left( \left\{ x \in \Omega : \limsup_{r \rightarrow 0} \frac{1}{r^{n-2}} \int_{B(x,r)} |u(y)| dy > 0 \right\} \right) = 0.$$

See [EG, p. 141 and p. 77].

By the above discussion, and by (2.1), there is a point  $x \in B_f$  such that (2.2) holds. We shall show that this is impossible. The method is a standard blow-up and normal families argument, used before in the study of branch sets, cf. [Re, II.§10], [MRV], [GMRV].

Without loss of generality, we assume that  $x = 0 = f(x)$ . Consider mappings

$$\begin{aligned} f_r &: \mathbf{B}^n \rightarrow \mathbf{R}^n, \\ f_r(x) &= r^{-1} f(rx), \quad r > 0, \end{aligned}$$

where  $\mathbf{B}^n$  is the open unit ball in  $\mathbf{R}^n$ . Because  $f$  is locally uniformly Lipschitz, there is a constant  $L > 0$  such that

$$f_r(\mathbf{B}^n) \subset B(0, L)$$

for all  $r > 0$  small enough. On the other hand, one also has that

$$B(0, 1/L) \subset f_r(\mathbf{B}^n)$$

by standard properties of *BLD*-mappings ([MV, Lemma 4.6]). The mappings  $f_r$  are uniformly *BLD* for all small  $r > 0$ , and it follows that for a subsequence  $f_k = f_{r_k}$ , the limit

$$\lim_{k \rightarrow \infty} f_k = F$$

defines a *BLD*-mapping,  $F: \mathbf{B}^n \rightarrow B(0, L)$ , such that  $B(0, 1/L) \subset F(\mathbf{B}^n)$  ([MV, Theorem 4.7]). Because the convergence of the sequence  $f_k$  is locally uniform, it follows from the basic degree theory that  $0 \in B_F$ ; that is,  $F$  is not a local homeomorphism at 0.

Next,

$$\int_{B(0, r_k)} |df(y) - (df)_{0, r_k}| dy = \int_{\mathbf{B}^n} |df_k(y) - (df_k)_{0,1}| dy \rightarrow 0$$

by assumption. By passing to another subsequence, we may assume that

$$(df_k)_{0,1} \rightarrow M$$

as  $k \rightarrow \infty$ , where  $M$  is an invertible  $n \times n$ -matrix and that

$$df_k(y) \rightarrow M \text{ at a.e. } y \in \mathbf{B}^n,$$

as  $k \rightarrow \infty$ . Note that all the matrices in question lie in a compact set of  $n \times n$ -matrices with definite positive distance from the zero locus of the determinant function.

Because  $f_k \rightarrow F$  uniformly and  $df_k \rightarrow M$  a.e., we must have that

$$dF(y) = M \text{ for a.e. } y \in \mathbf{B}^n.$$

But this means that, up to a linear change of coordinates,  $F$  is a conformal mapping if  $n \geq 3$ , or an analytic function if  $n = 2$ . (This follows from the generalized Liouville theorem for  $n \geq 3$  [Re, II §5.9], and from Weyl's lemma if  $n = 2$ .) In either case  $F$  must be a local homeomorphism, contradicting the result that  $0 \in B_F$ . The proof of the theorem is therefore complete.

### 3. Remarks on the quasiregular case

The argument in Section 2 works for general quasiregular mappings with some limitations. We use the terminology of [Ri].

For a (nonconstant) quasiregular map  $f$  in  $\mathbf{R}^n$  it is natural to consider the *dilatation tensor*

$$G_f(x) = \det df(x)^{-2/n} df(x)^* df(x),$$

defined almost everywhere in the domain of  $f$ . Thus,  $G_f$  is a bounded matrix valued measurable function with  $\det G_f(x) = 1$  almost everywhere. As in Section 2, we find that if  $G_f \in W^{1,p}$  for some  $1 \leq p \leq n$ , then

$$(3.1) \quad \lim_{r \rightarrow 0} \int_{B(x,r)} |G_f(y) - (G_f)_{x,r}| dy = 0$$

for  $x$  outside an exceptional set of Hausdorff  $(n-p)$ -measure zero. By a recent result of Martio et. al. [MRV, Lemma 3.2],  $f$  is locally invertible in a neighborhood of each point  $x$  such that (3.1) holds, provided  $n \geq 3$ . The proof in [MRV] is a blow-up and normal families argument similar to that in the previous section. (Note that  $G_f = Id$  for each holomorphic function  $f$  so that the dimensional restriction  $n \geq 3$  is necessary.) It follows in particular that every nonconstant quasiregular map of a domain in  $\mathbf{R}^n$ ,  $n \geq 3$ , is a local homeomorphism if  $G_f \in W^{1,n}$ . This can be improved somewhat:

**3.2. Proposition.** *Let  $f$  be a nonconstant quasiregular map of a domain in  $\mathbf{R}^n$ ,  $n \geq 3$ , with a nonempty branch set  $B_f$ . If  $f$  is Hölder continuous of order  $\alpha$  on  $B_f$  and if  $G_f$  belongs to the Sobolev space  $W^{1,p}$ , then  $p < n - \alpha(n-2)$ .*

*Proof.* Because  $\mathcal{H}_{n-2}(f(B_f)) > 0$  [Ri, III.5.3], the  $\alpha$ -Hölder continuity of  $f$  on  $B_f$  implies that  $\mathcal{H}_{\alpha(n-2)}(B_f) > 0$ , so that we must have  $n-p > \alpha(n-2)$  by the above discussion.  $\square$

Note that every quasiregular mapping  $f$  is locally Hölder continuous of order  $\alpha = K_I^{1/(1-n)}$ , where  $K_I = K_I(f)$  is the *inner dilatation* of  $f$  ([Ri, I.2.1, III.1.11]). Thus Proposition 3.2 is never vacuous.

If  $n = 3$ , then  $B_f = \emptyset$  or  $\mathcal{H}_1(B_f) > 0$  for all discrete and open maps, in particular for nonconstant quasiregular maps, by [MR, 2.20]. Moreover, if  $K_I(f) < 2$ , then  $f$  is locally Lipschitz continuous on  $B_f$  by a theorem of Martio [Ri, III.4.7]. We thus have the following result:

**3.3. Theorem.** *Let  $f$  be a nonconstant quasiregular map of a domain in  $\mathbf{R}^n$  with dilatation tensor  $G_f$  in the Sobolev space  $W^{1,2}$ . If either  $n = 3$ , or  $n \geq 4$  and the inner dilatation  $K_I$  of  $f$  is less than 2, then  $f$  is locally invertible.*

For  $n = 3$  Theorem 3.3 improves earlier results of Iwaniec [I], Manfredi [M], Gutlyanskii et. al. [GMRV], and Martio et. al. [MRV]. Because the dilatation tensor of the winding map belongs (locally) to  $W^{1,p}$  for all  $p < 2$ , it is tempting to believe that each nonconstant quasiregular map  $f$  with  $G_f \in W^{1,2}$  is locally invertible in all dimensions  $n \geq 3$ .

## REFERENCES

- [EG] Evans, L. C. and R. F. Gariepy, R. F., *Measure theory and fine properties of functions*, Studies in Advanced Mathematics. CRC Press, Boca Raton, FL, 1992.
- [GMRV] Gutlyanskii, V. Ya., O. Martio, V. Ryazanov, and M. Vuorinen, *On the local injectivity and asymptotic linearity of quasiregular mappings*, preprint Univ. Helsinki **122** (1997), (<ftp://www.math.helsinki.fi/pub/Reports/Preprint122.ps>).
- [HKM] Heinonen, J., T. Kilpeläinen, and O. Martio, *Nonlinear potential theory of degenerate elliptic equations*, Oxford University Press, Oxford, 1993.
- [HR] Heinonen, J. and S. Rickman, *Geometric branched covers between generalized manifolds*, (in preparation).
- [I] Iwaniec, T., *Regularity theorems for solutions of partial differential equations for quasiconformal mappings in several dimensions*, *Dissertationes Mathematicae CXCVIII* (1982), 1–48.
- [M] Manfredi, J. J., *Regularity of minima of functions with  $p$ -growth*, *J. Diff. Eq.* **76** (1988), 203–212.
- [MR] Martio, O. and S. Rickman, *Measure properties of the branch set and its image of quasiregular mappings*, *Ann. Acad. Sci Fenn Ser A.I. Math.* **541** (1973), 1–16.
- [MRV] Martio, O., V. Ryazanov, and M. Vuorinen, *BMO and injectivity of space quasiregular mappings*, preprint Univ. Helsinki **130** (1997), (<ftp://www.math.helsinki.fi/pub/Reports/Preprint130.ps>).
- [MV] Martio, O. and J. Väisälä, *Elliptic equations and maps of bounded length distortion*, *Math. Ann.* **282** (1988), 423–443.
- [Re] Reshetnyak, Yu. G., *Space mappings with bounded distortion*, *Translations of Mathematical Monographs 73*. American Mathematical Society, Providence, RI, 1989.
- [Ri] Rickman, S., *Quasiregular mappings*, Springer-Verlag, Berlin, 1993.
- [S] Sullivan, D., *Exterior  $d$ , the local degree, and smoothability*; In *Prospects of topology*, F. Quinn, Ed., Princeton University Press, Princeton, 1995.

Department of Mathematics  
 University of Michigan  
 Ann Arbor, MI 48109  
 U.S.A.  
 e-mail: juha@math.lsa.umich.edu

Department of Mathematics  
 University of Jyväskylä  
 P.O. Box 35 (Mattilanniemi D)  
 40351 Jyväskylä, Finland  
 e-mail: terok@math.jyu.fi